

SAINT-PETERSBURG UNIVERSITY

On the rights of the manuscript

Nassar

Ameed

CLINICAL AND BIOMETRIC ASSESSMENT OF THE SEVERITY OF THE
PHYSIOLOGICAL STATUS AND INJURIES IN SURVIVORS OF
CONCOFITANT MIDFACE TRAUMA

Scientific specialty: 3.1.2. Maxillofacial surgery

Thesis

for an academic degree
of Candidate of Medical Sciences

Translation from Russian

Research supervisor:

Madai Dmitrii Yur'evich

Doctor of Medical Sciences

Saint-Petersburg, 2024

Table of contents

LIST OF ABBREVIATIONS USED.....	4
INTRODUCTION.....	5
CHAPTER 1. CURRENT VIEWS ON THE PROBLEM OF ASSESSING THE SEVERITY OF PHYSIOLOGICAL STATUS AND INJURY IN SURVIVORS OF CONCOMITANT MIDFACE TRAUMA (LITERATURE REVIEW).....	14
1.1. Signs of Dysfunction of Vital Organs and Body Systems in Survivors of Concomitant Trauma.....	14
1.2. Objective Assessment of the Severity and Prediction of the Immediate Outcomes of Injuries.....	20
1.2.1. Methods for Assessing Severity and Predicting the Immediate Outcome of Injuries.....	22
1.2.1.1. Anatomical Methods for Assessing the Severity of Injuries.....	23
1.2.1.2. Physiological Methods for Assessing the Severity of Injuries.....	25
1.2.1.3. Combined Methods for Assessing the Severity of Injuries.....	28
1.2.2. Analysis of the Validity of Methods for Assessing Severity and Prognosis of Immediate Trauma Outcomes.....	31
1.3. Current Status of the Problem of Treating Survivors with Concomitant Midface Trauma.....	34
CHAPTER 2. MATERIALS AND METHODS OF RESEARCH.....	36
2.1. Research Materials.....	36
2.2. Research Methods.....	41
2.2.1. Methodology of Data Collection.....	41
2.2.2. Radiation Methods of Examination of Patients.....	42
2.2.3. Study of Clinical and Laboratory Signs Reflecting the Severity of the Physiological status of the Patients.....	43
2.2.4. Methods for Assessing the Severity of Concomitant Midface Trauma....	44
2.2.4.1. Injury Severity Score MFS-I (MT).....	44
2.2.4.2. Injury Severity Score ISS.....	45

2.2.5. Statistical Processing of Research Data.....	46
CHAPTER 3. EXPLORATORY ANALYSIS OF RESEARCH PAPER DATA.....	48
3.1. Exploratory Analysis of the Research Material.....	48
3.2. Exploratory Analysis of Studied Physiological Signs.....	49
CHAPTER 4. CLINICAL AND BIOMETRIC APPROACH TO ASSESSING THE SEVERITY OF THE PHYSIOLOGICAL STATUS OF PATIENTS WITH CONCOMITANT MIDFACE TRAUMA.....	59
CHAPTER 5. METHODOLOGY FOR PREDICTING THE IMMEDIATE OUTCOMES OF COMCOMITANT MIDFACE TRAUMA	75
5.1. The First Variant of the Discriminant Function.....	77
5.2. The Second Variant of the Discriminant Function.....	79
CHAPTER 6. TREATMENT OF PATIENTS WITH CONCOMITANT MIDFACE TRAUMA	82
6.1. Tactics of Multi-Stage Surgical Treatment of Patients with Concomitant Midface Trauma.....	82
6.2. Treatment of Patients with Severe Concomitant Midface Trauma	85
6.2.1. Clinical Characteristics of Patients with Severe Concomitant Midface Trauma.....	85
6.2.2. Results of Treatment of Patients with Severe Concomitant Midface Trauma	88
6.3. Treatment of Patients with Extremely Severe Concomitant Midface Trauma.....	94
6.3.1. Clinical Characteristics of Patients with Extremely Severe Concomitant Midface Trauma.....	94
6.3.2. Results of Treatment of Patients with Extremely Severe Concomitant Midface Trauma	97
CONCLUSION.....	104
FINDINGS.....	110
PRACTICAL RECOMMENDATIONS.....	112
REFERENCES.....	113

LIST OF ABBREVIATIONS USED

TBI –traumatic brain injury;

MFR –maxillofacial region;

MZF –middle zone of the face;

ALV –artificial lung ventilation;

–ICU Intensive Care Unit;

MFS –military field surgery;

MST –tactics of multi-stage surgical treatment;

TD –traumatic disease;

HR– heart rate;

INR– International Normalized Ratio;

pH –hydrogen index;

FiO₂– fraction of oxygen on inspiration;

PO₂ –partial pressure of oxygen in arterial blood;

Na⁺– sodium ions;

K⁺ –potassium ions;

GCS – Glasgow Coma Scale;

ISS – Injury Severity Score;

AIS –Abbreviated Injury Scale;

RTS –Revised Trauma Score;

APACHE –Acute Physiology and Chronic Health Evaluation;

ICISS-9– International Classification of Diseases (ICD-9) based Injury Severity Score;

SAPS –Simplified Acute Physiology Score;

SOFA –Sequential Organ Failure Assessment;

TRISS –Trauma and Injury Severity Score;

RISC –Revised Injury Severity Classification;

TMPM is a –Trauma Mortality Prediction Model.

INTRODUCTION

Relevance of the study

The problem of severe concomitant trauma in medicine remains as relevant as ever, despite the active development and introduction of modern technologies in the diagnosis and treatment of patients with traumatic injuries in recent decades [7, 8, 15, 17].

This is due, firstly, to the wide prevalence of various types of injuries in modern society, which is associated with growing urbanisation, accompanied by the wide availability of means of transportation and personal protection [5, 7, 23, 55]. Secondly, to a variety of clinical manifestations of concomitant injuries, the high probability of developing complications in the patients, and the high risk of an unfavourable outcome compared with an isolated injury [45, 48, 49]. Thirdly, traumatism has economic and social consequences due to the fact that the majority of patients are represented by men of working age (20-50 years) [132, 151, 159, 162, 168, 185, 219], and also with high financial costs of treatment and rehabilitation of patients [87, 139].

Among the causes of death of the population in peacetime, concomitant trauma ranks third after oncological diseases and diseases of the cardiovascular system [21, 27, 86, 91, 134, 213] and it is the main cause of mortality of young and able-bodied people [23, 63, 153].

Concomitant craniofacial trauma is among the most severe and life-threatening injuries and is characterised by a violation of the central mechanisms of regulation of various organs and systems of the body [28].

The facial skeleton is usually divided into three anatomical regions: upper, middle and lower. The upper border of the middle zone of the face (MZF) is marked by a line drawn from the zygomatic suture on one side through the upper ocular edges, the left maxillary sutures, and nasolabial sutures to the zygomatic suture on the other side. The lower border of the MZF is located at the level of the occlusal plane of the

teeth of the upper jaw or the alveolar edge, in the case of their absence [40, 90, 100]. The posterior sections of the MFZ are formed by the pterygoid processes of the sphenoid bone [40]. MFZ is composed of the following bones and bone structures: maxillae, zygomatic bones, palatine bones, nasal bones, lacrimal bones, inferior nasal concha, vomer [40, 54, 72], zygomatic processes of the temporal bones, ethmoid bone, and wing processes of the sphenoid bone [40, 72].

MFZ has a complex anatomical structure [29, 73, 82, 100, 158, 189]. The presence of visual analysers, ENT organs, and anatomical proximity to the brain, as well as frequent combination with traumatic brain injury (TBI) [16, 54, 78, 82, 107, 222], frequently cause functional and aesthetic complications in the midface trauma [14, 53, 54]. According to a group of researchers, TBI is combined with midface injuries in 6-30% of cases [44, 94, 99].

The modern concept of treatment of patients with concomitant trauma can be characterised as pathogenetically justified. According to this concept, the extent of medical care required is determined according to the nature of the pathogenetic processes occurring in the injured person after injury and reflecting the severity of his or her physiological status [128].

Thus, assessment of trauma severity is a high priority for modern therapeutic and diagnostic management tactics. This is the basis for deciding on the extent, nature, and order of medical care [31].

The degree of exploration of the research topic

The methods for assessing the severity of injuries have been actively developing since the 70s of the last century. To date, more than 50 scales and methods have been developed to solve this problem. Most of these methods have not survived the test of time. In the last decade, the attention of researchers has been directed to the development of specific methods for assessing the severity of injuries for a specific type of injury or damage to certain organs and anatomical areas.

The main objective of the researchers who studied the problem of midface injury was to develop methods of diagnosis and treatment in order to eliminate local aesthetic and functional impairment after injury and to improve the treatment outcomes of patients with isolated midface trauma.

The authors studying the problems of concomitant midface trauma investigated the timing of reconstructive operations of damaged structures of the maxillofacial region (MFR), the rationale for the tactics of MST taking into account the anatomical structure of the craniofacial skeleton and the effectiveness of the MST tactics in patients with a severe concomitant trauma of the MFR as part of polytrauma [10, 25, 28, 51]. The results of MST tactics in patients with extremely severe concomitant midface trauma as a separate group have not been studied.

A review of the available literature showed that there have been no scientific studies on the problem of assessing the severity of the physiological status in patients of concomitant midface trauma. There is also a lack of scientific work on the study of signs of dysfunction of vital organs and body systems in the dynamics of the course of traumatic disease, which is extremely important for understanding its pathogenesis and building tactics for multi-stage surgical treatment of patients with this type of injury.

Given the high mortality rate, the risk of infectious and noninfectious, including life-threatening complications, and the increasing number of patients with concomitant midface injuries, this problem requires further study to justify the choice of surgical treatment tactics on an objective basis, taking into account the severity of the physiological status of the patients.

The purpose of the study

To conduct a study of clinical and laboratory signs for an objective assessment of the severity of the physiological status of the patients and prediction of the immediate outcomes of concomitant midface trauma.

Objectives of the study

1. To search for syndrome complexes and signs characterising the severity of the physiological status of patients with concomitant midface trauma in the dynamics of the course of traumatic disease.
2. To develop a methodology for predicting the immediate outcomes of concomitant midface trauma.
3. To evaluate the effectiveness of tactics of multi-stage surgical treatment in patients with severe and extremely severe concomitant midface trauma.

Scientific novelty

In the present study, the clinical and laboratory parameters characterising the severity of the physiological status in survivors of concomitant midface trauma in the dynamics of the course of traumatic disease were studied for the first time. It has been proved that the nature of morphological relations of the main cells of the immune system and the severity of general intoxication and infectious complications determine the course of the traumatic disease in dynamics.

The third post-injury day was found to be the most optimal for predicting the immediate outcome of a concomitant midface trauma. A methodology for predicting the immediate outcome of a concomitant midface trauma has been developed and presented for the first time.

A surgical care algorithm has been established for survivors of concomitant midface trauma. The results of multistage surgical treatment tactics on multiple survivors with extremely severe concomitant midface trauma were studied for the first time. This has been shown to reduce the likelihood of death and infectious and non-infectious complications by taking into account the severity of the overall physiological status of the injured person.

Theoretical and practical significance of the work

The main stages of pathogenesis of traumatic disease in patients with concomitant midface trauma have been investigated. Some clinical and laboratory indicators that accurately reflect the severity of functional disruptions of the patients' critical organs and systems in the dynamics of the course of the traumatic disease have been studied.

The significance of the signs characterising morphological and cooperative relations of the main cells of the immune system and indicators of the severity of general intoxication of the organism for evaluation of the severity of the general state of patients in dynamics has been evaluated. The correlations and the level of dependency of these attributes have been revealed.

A comparative biometric analysis algorithm using two-time directions (time course of traumatic disease and time to outcome) and factor analysis based on the principle of shifting all realisations to the same time point with similar clinical significance for the patients studied, followed by a comparative analysis of the identified factors in the dynamics of the course of traumatic disease has been developed.

A rational algorithm for the treatment of patients with extremely severe concomitant midface trauma using the tactics of multistage surgical treatment and taking into account the severity of the physiological status, which helped to reduce mortality by 17.9% and the incidence of infectious complications by 9.2%, has been established.

The object of research: the man.

Subject of the study: clinical and laboratory physiological signs reflecting the functional state of vital organs and body systems of the patients, spiral computer tomograms and head X-rays, and case histories of patients with concomitant midface trauma.

Methodology and methods of research

The goal of the work was to investigate clinical and laboratory signs to objectively assess the severity of the physiological status of the injured and to predict the immediate outcome of a concomitant midface trauma.

The research methods were based on the measurements of clinical and laboratory indexes, characterising the severity of the general state of the injured, a comparative biometric analysis of the values of these signs among the injured, depending on the nearest outcomes of the concomitant trauma in order to determine their significance in estimating the severity of the general state of the injured in dynamics, the course of the traumatic disease. and prognosis of the nearest outcomes of the concomitant trauma. The work uses laboratory, clinical, and instrumental research methods.

The main scientific results

1. The dynamics of the average values of the leukocyte intoxication index, the neutrophil-lymphocyte ratio, and the absolute number of lymphocytes and monocytes reflects the course of the traumatic disease and the severity of the physiological status of patients with a concomitant midface trauma. LII is an independent objective criterion reflecting the severity of the physiological status of the patients with a concomitant midface trauma and the severity of the course of the traumatic disease as a whole [35].

2. Taking into account the set of signs and time points studied by us, the third day after the occurrence of the injury is the most optimal point for predicting the immediate outcomes of the concomitant midface trauma. A method for predicting the immediate outcomes of a concomitant midface trauma has been developed. It has been established that when a patient experiences cerebral edema, the likelihood of lethal outcome increases significantly [36].

3. In the cohort of patients with a severe concomitant midface trauma the use of multi-stage surgical treatment tactic, taking into account the severity of the

physiological status of the patients, led to a decrease in the frequency of infectious complications by 27%, non-infectious complications - by 5.8% and a decrease in mortality by 19%; in the cohort of patients with an extremely severe concomitant midface trauma - to a decrease in the frequency of infectious complications by 9.2%, non-infectious complications by 25.4% and a decrease in mortality by 17.9% [64].

The main provisions submitted for defence

1. An objective method of assessing the severity of injuries and the physiological status of patients with a concomitant midface trauma is a mandatory tool in determining the scope of permissible and necessary surgical care and the timing of its provision.

2. The nature of morphological ratios of the main cells of the immune system, the severity of general intoxication of the body and infectious complications reflect the nature of the course of the traumatic disease.

3. The leukocyte intoxication index is an objective criterion for assessing the severity of the physiological status of the patients in the dynamics of the course of traumatic disease.

4. The best treatment results are achieved by applying rational tactics of multi-stage surgical treatment with an objective dynamic assessment of the severity of their physiological status.

The degree of validity of the results of the study

In the course of the dissertation research, original methods and methods of collecting and subsequent processing of primary information were used in order to form a preliminary dataset for biometric analysis. The reliability of scientific statements, conclusions, and practical recommendations is ensured by compliance with the methodological systemic approach at all stages of the study and the use of adequate and modern methods of statistical data processing. Based on the collected clinical

material, the issues of assessing the severity of the physiological status of the patients were studied mainly on the basis of objective laboratory indicators, which made it possible to develop and implement an algorithm for providing surgical care to patients with severe and extremely severe concomitant midface trauma.

Implementation of the results

The results of the dissertation research have been implemented in the practical activities of the First Neurosurgical Department (for patients with combined craniofacial trauma and damage to the organs of hearing and vision) of St Petersburg State Medical Institution 'Alexandrovskaya Hospital' and St Petersburg State Medical Institution 'Dzhanelidze Institute of Emergency Medicine'.

The results of the study have been used in the form of a lecture and are used in the educational process by the Department of Maxillofacial Surgery and Surgical Dentistry of St Petersburg University.

Approbation of the work

The main provisions of this dissertation were discussed at a meeting of the Department of Maxillofacial Surgery and Surgical Dentistry of St Petersburg University. The results of the work were reported at scientific conferences: the 5th All-Russian Congress with international participation 'Medical Care for Injuries. New in organisation and technology. Prospects of Import Substitution in Russia' (St Petersburg, 2020); All-Russian Conference on Natural Sciences and Humanities with international participation 'Science of St Petersburg University - 2020' (St Petersburg, 2020); All-Russian scientific and practical conference dedicated to the 100th anniversary of the birth of Professor V.A. Malyshev 'Topical Issues of Maxillofacial Surgery and Dentistry' (St Petersburg, 2022); All-Russian Interdepartmental Scientific and Practical Conference 'Modern Approaches to the Diagnosis and Treatment of Surgical Diseases, Injuries, and their Complications' (Moscow, 2022).

Publications

11 research papers have been published on the materials of the study, including 5 articles in peer-reviewed scientific journals recommended by the Higher Attestation Commission of the Russian Federation.

Personal contribution of the author

The author independently analysed the works of foreign and domestic authors, and defined the goal, objectives, methodology, design, and stages of the dissertation research. The researcher has developed and implemented an algorithm for collecting primary data and forming a preliminary representative dataset. The author was directly involved in the mathematical and statistical analysis of the research data. The author participated in the implementation of a complex of clinical, laboratory, and instrumental methods of examination.

In his co-authored scientific papers, the researcher has provided the main theoretical background for problem definition, design, clinical material collection, and drawing conclusions.

Scope and structure of the thesis

The thesis consists of 142 typewritten pages, including an introduction, a literature review, a chapter of materials and methods, a chapter of exploratory analysis, three chapters of own research, a conclusion, findings, practical recommendations, and a list of references including 100 Russian and 122 foreign sources.

CHAPTER 1.

CURRENT VIEWS ON THE PROBLEM OF ASSESSING THE SEVERITY OF PHYSIOLOGICAL STATUS AND INJURY IN SURVIVORS OF CONCOMITANT MIDFACE TRAUMA (LITERATURE REVIEW).

1.1. Signs of Dysfunction of Vital Organs and Body Systems in Survivors of Concomitant Trauma

Special attention is paid to the search for signs indicating the severity of the violation of vital functions of the body when developing methods for assessing the severity of injury [33]. The higher the specificity of these signs to a certain nosology or damage, the higher the accuracy of the developed method. In the case of the concomitant midface trauma, the analysis of the special literature shows insufficient exploration of this topic [79].

The most common parameters used in assessing the severity of the physiological status of the patients is the level of blood pressure. According to A.N. Kolesnikov and other authors, the systolic blood pressure is a significant clinical parameter reflecting the degree of post-traumatic haemodynamic disorders and the likelihood of shock, especially at the prehospital stage [38], regardless of the localisation of the dominant injury [24, 32, 145, 172, 173, 182, 198].

According to S.D. Saverio et al., hypotension on admission to hospital with a systolic blood pressure of 90-100 mmHg against a background of blood loss and traumatic shock, with an Injury Severity Score of >16, can be considered a factor of adverse outcome [119].

Yu.V.Puras et al. similarly interpret episodes of arterial hypotension as an adverse outcome factor in patients with concomitant trauma [76]. Others have suggested that a rise in systolic blood pressure above 160 mmHg on admission to hospital has a high predictive value as a prognostic factor for death [200].

Another indicator of haemodynamics, often used in assessing the severity of the physiological status, is the heart rate (HR). E.J. Ley et al. found that the risk of death

in patients increases with a heart rate of ≤ 70 or ≥ 89 beats per minute at the time of admission to the clinic. The highest risk of death is observed with heart rate at the time of admission of <60 or ≥ 100 beats per minute [109].

Some researchers believe that heart rate alone is not a significant indicator of the severity of the patient's physiological status and the need for emergency surgical interventions when the patient is admitted to hospital [142].

G.P. Victorino et al. established that the relationship between heart rate and systolic blood pressure at the time of admission of the patient to hospital is not significant, and tachycardia is not a reliable sign of hypotension [218].

The haemoglobin level is a significant sign in assessing the severity of the physiological status of patients with severe blood loss. S.M. Alamshah et al. interpret a low haemoglobin level on admission of the patient to hospital as a valuable prognostic factor of an unfavourable outcome of a severe injury [135].

J.H. Holstein, et al. defined the haemoglobin level of 67 ± 29 g/l, the systolic blood pressure of 77 ± 27 mmHg, and the severity of the injury on the ISS scale at 35 ± 16 points on admission as factors of an unfavourable outcome for patients with severe concomitant trauma [145].

S. Majercik et al. indicate the importance of the red blood cell distribution index - RDW (Red Cell Distribution Width) in predicting the lethal outcome of a severe injury. The index, according to the authors, makes it possible to make an accurate prediction of the 30-day mortality in male patients, and the annual mortality in male and female patients [181].

The researchers discuss the significance of blood gas composition indicators in assessing the severity of the physiological status of patients with concomitant trauma. According to T.M. Dumont et al., an increase in the partial pressure of blood carbon dioxide (pCO_2) >45 mmHg (hypercapnia) or its decrease <35 mmHg (hypocapnia) in patients with severe concomitant trauma at the time of admission to hospital increases the probability of death [150]. According to other data, the gas composition of blood is not a significant factor in predicting the fatal outcome in patients with concomitant trauma [143].

B. indicate no correlation between the Glasgow Coma Scale (GCS) score and blood gas values (pH, pO₂, pCO₂, HCO₃⁻ bicarbonate and SO₂ saturation) during the first 4 hours after admission in a patient with severe concomitant trauma. According to the authors, there was also no significant difference in the average values of blood gas composition in patients with favourable and unfavourable outcomes [131].

Redistributive leucocytosis often occurs after injury and is associated with significant physiological stress. It occurs as a result of the demargination of leukocytes from the walls of blood vessels under the action of catecholamines [124, 156, 179].

When examining the number of leukocytes in the patients at the time of admission to hospital, D.C. Chang et al. found that there was a connection between the severity of injuries on the ISS >15 scale and an increase in the number of leukocytes to $11,9 \times 10^9/l$, as well as between the sum of scores on the scale of GCS ≤ 8 and leukocytosis at the level of $13.0 \times 10^9/l$ [136].

Claudia A. Santucci et al. when comparing the number of leukocytes in patients with severe trauma and in patients with mild and moderate trauma at the time of admission to hospital, found that the number of leukocytes in the two groups significantly differed. The authors established weak direct correlation (coefficient to correlation = 0.37) of the severity of injuries on the ISS scale and the number of leukocytes on admission [187].

In the study by S. Lam et al., comparing the total number of leukocytes and their differential counting, as well as the size of blood neutrophils in patients with favourable and lethal outcomes within seven days after injury, found that the size of neutrophils upon admission to hospital is a significant prognostic factor of seven-day mortality. The total number of leukocytes and their differential counting had no prognostic value [124].

There is some literature indicating the importance of neutrophil to lymphocyte ratios, measured at the time of admission, in predicting adverse outcome of concomitant trauma [194], including concomitant TBI [164].

To describe coagulopathy, hypothermia, and metabolic acidosis, the term 'triad of death' is often used in foreign scientific literature. These signs are associated with

impaired tissue perfusion, haemostatic disorders, and the development of multiple organ failure [74, 133].

According to M.G. Balverde et al., the 'triad of death' syndrome can be classified as an independent adverse outcome factor for combined trauma, and is characteristic of injuries with an ISS score of >30-35. The 'triad of death' syndrome includes coagulopathy (international normalized ratio (INR)>1.5), hypothermia (temperature <35 ° C) and metabolic acidosis (pH <7.2) [177].

B. Mitra et al. found that the probability of a fatal outcome in patients with the 'triad of death' is 48%, and with INR >3.2 - 100% [210].

Most researchers determine the level of coagulopathy, which can be used as an independent factor of an unfavourable outcome of a concomitant injury, by the INR indicator. According to M.E. Kutcher et al., this indicator should exceed 1.3 [113], and according to other data - 1.5 [110, 121].

M. Kapan and other researchers have identified hypothermia below 35° on admission to the hospital, as an independent factor of the unfavourable outcome of a concomitant injury [137, 198, 201]. According to other researchers, hypothermia cannot be taken as an independent factor of an unfavourable outcome of a concomitant injury [137, 190], since it is only a manifestation of other post-traumatic disorders, such as coagulopathy, acidosis, or severe blood loss [138].

Blood lactate concentration reflects the level of oxygen deprivation of tissues, acidosis, and the acid-base state of the body. Hyperlactemia is one of the prognostic factors of an unfavourable outcome of a concomitant injury [175, 204].

Normally, the blood lactate level does not exceed 1 mmol/L. The lowest level of hyperlactemia, which can be taken as an indicator of oxygen deprivation of tissues, acidosis, and severe disorders of the acid-base state of the body, is 2 mmol/L [174], or according to other data 4.1 mmol/L [74].

S.M. Alamshah et al. defined metabolic acidosis with a blood pH <7.2 was determined as an independent factor of an unfavourable outcome of injury [135]. M.T. Gokdemir et al. Gokdemir et al. note a correlation between total plasma oxidative status in severely injured patients and mortality and injury severity on the ISS and Revised

Trauma Scale - RTS. The authors suggest using this parameter as an early biomarker of oxidative stress to control the severity of patients with multiple trauma in its acute period [184].

According to E. Fitzsullivan et al., the level of blood plasma bicarbonate HCO_3^- has a strong positive correlation with the level of plasma base deficiency, both at the time of admission hospital ($r = 0.80$) and at the entire stage of hospital treatment ($r = 0.85$). Both indicators more accurately reflect the level of metabolic acidosis than the pH or blood lactate level. The authors also note the presence of a significant difference in the average indicators of deficiency of bases and bicarbonate HCO_3^- in patients with favourable and unfavourable outcomes (2.5 and 5.2; 17.7 and 19.8 mmol/L, respectively) [188].

Plasma potassium and sodium levels are often used to describe and assess water-electrolyte imbalance. In their study, V. Morell et al. found that hypokalemia with a potassium level (K^+) < 3.6 mmol/L on admission correlated with the severity of injuries on the ISS scale and the duration of hospital stay of the patients [160].

Investigation of potassium (K^+) levels on admission in patients with concomitant trauma in the work of A.L. Beal et al. showed that hypokalaemia (< 3.6 mmol/L) was more common in patients with TBI. In patients with hypokalemia, the sum of points on the GCS was lower, and the sum of points on the ISS scale was higher than in patients with normal potassium levels. The need for artificial lung ventilation (ALV), the duration of stay in the intensive care unit (ICU) and the duration of hospital stay were greater in patients with hypokalemia. The authors also noted that the sum of the scores on GCS in patients with potassium levels < 3.1 mmol/L was lower than in patients with potassium levels in the range of 3.1-3.6 mmol/L [147].

S. Pomeranz et al., when comparing the level of potassium on admission to hospital in the group of patients with isolated severe TBI (≤ 7 points according to GCS) and the group of patients with concomitant trauma, but without TBI, found that the level of blood potassium in the patients of the first group was significantly lower than in the patients of the second group. The average potassium level in the patients of the first and second groups was 3.1 ± 0.4 mmol/L and 3.5 ± 1.1 mmol/L, respectively.

According to the authors, this difference is due to large catecholamine emissions in severe TBI with subsequent stimulation of sodium-potassium pumps [169].

A. Vedantam et al. when studying the blood sodium level (Na^+) in the first week after admission in patients with severe TBI, found that hypernatremia ($\text{Na}^+ >150$ mmol/L) is associated with the likelihood of acute renal failure and death [217]. Other researchers have also found that hypernatremia ($\text{Na}^+ = 148-152$ mmol/L) in patients with severe TBI (≤ 8 points according to GCS) is associated with an increased probability of an unfavourable outcome of the injury [205]. Similar results were obtained by Hoffman H. et al., who found that hypernatremia in patients with severe TBI is associated with an increase in the probability of death and the duration of hospital stay [144, 146].

In a study of factors associated with lethal outcome in patients with grade III traumatic shock, Stukanov et al. found the following parameters to be associated: hypothermia (temperature $<35.7^\circ\text{C}$), increased lactate concentration in venous blood to 4.1 mmol/L, venous blood pH to 7.19, ionised calcium in venous blood to 0.3 mmol/L and activated partial thromboplastin time to 59 sec. [74].

According to G.A. Alexiou, an increase in blood sugar levels is often found in patients with concomitant trauma [115]. J. Kreutziger et al. found that the blood sugar level on admission to hospital of >10 mmol/L or <2.8 is more common in patients with traumatic shock [114].

In various publications, researchers have assessed the importance of blood sugar levels in predicting the outcome of injury. According to some researchers, the blood sugar level >8.8 mmol/L is a significant prognostic factor of an unfavourable outcome of TBI [183]. According to other data, blood glucose level is not a significant factor in predicting an unfavourable outcome of TBI [113].

F. Salehpour et al. indicate that the blood sugar level is not a significant factor in predicting an unfavourable outcome of severe TBI. The difference in the average values of blood glucose levels in patients with favourable and unfavourable outcomes was not significant (10.3 ± 2.9 and 10.4 ± 4.2 mmol/L, respectively). The authors also

did not reveal a significant correlation between glucose levels and the sum of scores on the GCS [118].

Analysing the above research results of foreign and domestic authors, we can talk about a variety of signs of post-traumatic disorders of the functions of organs and body systems, which is associated with the complexity and nature of their interaction in response to traumatic effects.

It should be emphasised that most researchers have studied these signs in trauma patients, irrespective of their features and localisation, including the location of the leading injury in terms of severity. Most of these studies were carried out at a specific time point (most often at the time of admission to the clinic), or at a short time interval (up to 3 days).

Thus, an important and, at the same time, unsolved problem of an objective assessment of the severity of a concomitant midface trauma is the issue of identifying physiological signs reflecting the severity of the physiological status of the patients in the the course of traumatic disease (TB).

1.2. Objective Assessment of the Severity and Prediction of the Immediate Outcomes of Injuries

The severity of injury is a complex concept that includes the severity of injuries and the severity of the physiological status. The severity of the injuries is a relatively stable indicator reflecting the morphological component of the injury (the result of the interaction of morphological structures of the body with the damaging agent). It depends on the localisation, the extent of anatomical damage, and the functional significance of the damaged organ [20, 32, 47, 49, 50, 57, 65, 87, 92].

The severity of the physiological status reflects the functional component of the injury and depends on the severity of functional disorders, the time from the moment of injury, and the initial state and reactivity of the body. The severity of the physiological status is a labile indicator and it can change at different stages of medical care, depending on its scope and quality [20, 32, 47, 49, 50, 57, 65, 87, 92]. The severity

of the physiological status and the severity of the damage are interrelated, but not interchangeable.

Assessment of the severity of injuries serves to address the following tasks:

1. Classification of injuries;
2. Sorting of patients;
3. Predicting the outcome of injury and the likelihood of complications [75];
4. Determination of the composition of the team of specialists involved in the process of diagnosis and treatment, and the priority of providing assistance to patients [80];
5. Determination of the scope of diagnostic tests [81];
6. Comparison and analysis of the quality of medical care provided and identification of its shortcomings [81];
7. Optimisation of treatment tactics at different stages of medical care;
8. Standardisation and objectification of the approach to solving various tasks in the provision of medical care and reducing the likelihood of errors in diagnosis and treatment [81].

Prior to the development and implementation of modern systems of objective assessment of the severity of injuries in clinical practice, descriptive methods were traditionally used to indicate the severity of injury [30, 34].

Over the history of the study of this problem since the early 1970s, domestic and foreign specialists have developed and tested dozens of methods and scales [33, 62, 80]. A large number of proposed methods indicate the complexity and importance of this problem and the lack of a single and generally accepted opinion on its solution, both in world and domestic medicine [31].

The lack of a unified system for assessing the severity of injuries makes it difficult to compare the effectiveness of various methods of treating patients, including the level of mortality and the risk of complications with the same severity of injuries [62, 83, 84, 85, 170, 209].

The need for a universal and unified system for assessing the severity of injuries is obvious and does not cause disagreement among specialists. However, the

development and implementation of a unified method for assessing the severity of injury proved to be a difficult task.

The human body is a very complex system. The variety of injuries and disorders occurring in the body after injury make it difficult to create a universal method [79]. It is debatable which parameters most accurately reflect the severity of the physiological status of the patient and should be taken as the basis of a universal method.

The development of a unified method for assessing the severity of injuries is also hampered by differences in equipment in hospitals and constant changes in the care of the injured, due to the introduction of new technologies in the treatment process [81]. The accuracy of the methods developed in some institutions may decrease when they are used in other institutions with a lower level of quality of medical care [79].

1.2.1. Methods for Assessing Severity and Predicting the Immediate Outcome of Injuries

Depending on the characteristics of the parameters used in the assessment, the methods of assessing severity can be divided into 3 groups:

1. Anatomical methods - based on the assessment of the severity of morphological disorders of various tissues and organs.

2. Physiological methods - based on the assessment of physiological parameters reflecting the severity of the physiological status of the patient.

3. Combined methods - include assessment of both morphological disorders of tissues and organs, and the severity of the physiological status of the patient [52, 65, 75, 83, 98].

The methods for assessing the severity of injuries vary depending on the task they solve. They can also be divided into general and specific methods of assessing severity. The general methods apply to all types of injuries or diseases, regardless of their localisation and nature. Specific methods are used in the treatment of certain diseases or in case of damage to a certain organ or organs and tissues of one anatomical

area. A brief description of the most clinically known scales and methods for assessing the severity and prognosis of injury outcomes will be presented below.

1.2.1.1. Anatomical Methods for Assessing the Severity of Injuries

Among the first methods of assessing the severity of injuries was the Abbreviated Injury Scale - AIS, developed in the USA in 1969. The scale is based on a score assessment of the severity of the most critical injury [123, 148, 180]. Since that time, the scale has been revised and supplemented 8 times and the last one was in 2015 [106].

Since the developers of the AIS scale did not try to solve a medical problem, but sought to create a system for determining insurance payments depending on injuries sustained in car accidents, the AIS scale in its initial form was not adapted for use in clinical practice. Due to the fact that the AIS injury severity scale is scored on the basis of the number of points of one most severe injury, it is considered unsuitable for use with multiple and concomitant injuries [199]. Nevertheless, this scale laid the foundation for the development of modern methods for assessing the severity of injuries and the introduction of a point-based severity assessment instead of a descriptive one.

S.P. Baker et al. in 1974 proposed a scale of injury severity - ISS (Injury severity score), created on the basis of the AIS scale. In the ISS scale, the authors used the damage codes proposed in the AIS scale. The severity of injuries on the ISS scale includes the sum of the squares of the codes of the 3 most severe injuries in 3 different anatomical areas [202].

Thus, the sum of points on the ISS scale varies from 1 to 75 points. The authors proposed the following interpretation of the assessment: 1-8 points - light injuries; 9-15 points - moderate injuries; 16-24 - serious injuries with a high probability of survival; 25-49 - severe injuries with a high probability of death; 50-74 - critical injuries; 75 points – injuries incompatible with life. If an injury is coded at 6 on the

AIS scale, it is considered incompatible with life and automatically coded at 75 on the ISS scale [202].

The ISS is by far the most widely used and discussed scientific method for assessing injury severity. It is used as a standard for the classification of injuries in many countries, including the USA, Australia, and most European countries [196].

However, the ISS method has a number of disadvantages. P.A. Seliverstov et al. indicates that the ISS scale underestimates the importance of severe TBI in determining the outcome of a combined injury. The authors also note that the number of points assigned to each injury on this scale does not always correspond to their value in determining the outcome of the injury [79].

The next drawback of the ISS method is related to the method of assessing the severity of multiple injuries. The ISS scale takes into account only one most severe injury within one anatomical area. This means that if there are several severe injuries in one anatomical area, only the most significant one is taken into account, and the role of other injuries in determining the outcome of the injury is not evaluated. At the same time, other, lighter injuries occurring in other anatomical areas are assessed, even though their contribution to determining the outcome of the injury is clearly less [38, 165].

To address this problem, T. Osler et al. proposed using a new injury severity scale - NISS (New Injury Severity Score), which is essentially a modification of the ISS scale. When assessing the severity of injuries on this scale, the three most severe injuries are taken into account, regardless of their localisation [165].

Another modification of the ISS scale was presented by W.S. Copes et al. in 1990. Anatomic Profile Score - APS takes into account three injuries with a severity > of 3 points on the AIS scale. One injury in the head and spinal cord area, one in the chest and neck area, and one in other anatomical areas are taken into account. The calculation on this scale is carried out using the logistic regression equation [178]. Due to the complexity of the calculation, this scale is not widely used among specialists.

In an attempt to create a unified international scale for assessing the severity of injuries, T. Osler et al. developed a method for assessing the severity of injuries based

on the International Classification of Diseases 9 (ICD-9) - ICISS-9 (International Classification of Disease-9 (ICD-9) based Injury Severity Score). The severity of the injury on this scale is assessed using the SRR (survival risk ratios) survival coefficient calculated for each specific injury [148].

According to the authors, the accuracy of this method is due to the fact that it was developed on an empirical basis, with the inclusion of a large number of patients (in ICISS-9 = 300,000 patients) and with the calculation of the probability of a fatal outcome for each injury in a retrospective way [148].

The ability to calculate the survival rate for each injury reflected in the ICD-9 raises questions, since most often these injuries, including injuries leading to a lethal outcome, occur in practice in a combined form [79].

Due to the fact that the International Classification of Diseases (ICD) is periodically revised, as well as the fact that the quality of medical care is changing, this method requires periodic revision [81].

The level of medical care differs in different states and, accordingly, the probability of mortality for certain injuries may vary depending on the locality, which calls into question the accuracy and universality of this method.

T. Osler et al. in 2008 presented a model for predicting the probability of a fatal outcome of patients - TPM (Trauma Mortality Prediction Model). The model is based on the assessment of the 5 most severe injuries, regardless of their localisation, using the AIS scale followed by mathematical modeling [105].

We consider the Military Field Surgery (MFS) injury severity scale in the combined methods section, together with other MFS severity scales, as it is part of a combined severity assessment system.

1.2.1.2. Physiological Methods for Assessing the Severity of Injuries

Physiological methods for assessing the severity of the physiological status of the injured person are based on the following principle: the severity of the

physiological status of the injured person can be measured by the degree of deviation from the normal physiological indicators after injury.

To assess the severity of TBI and the level of consciousness of the patient, the Glasgow Coma Scale (GCS) is widely used. The parameters evaluated on this scale are: motor reaction, speech response, and eye opening. The number of points received varies in the range of 3-15. The higher the score, the less severe the trauma and the less pronounced the impairment of consciousness [33, 193].

Despite the high level of subjectivity of the assessment method on this scale, it has found wide application, due to its simplicity and convenience. Literature data indicate high levels of sensitivity (79-97%) and specificity (84-97%) of this scale, both for assessing the severity of TBI and for predicting the fatal outcome in severe TBI [2, 76, 172].

H.R. Champion et al. in 1980 presented a scale for sorting patients - TS (Triage Score). The sorting of the patients, according to this scale, is based on the assessment of the following 3 parameters: chest excursion during breathing, capillary filling, and the level of consciousness according to the scale. The scale allows for pre-hospital care to determine where to take the patient, to a specialised or general hospital [112].

In 1981, the first modified version of the scale for sorting patients was proposed under the name 'Trauma Score'. In addition to the level of consciousness in the GCS and the degree of capillary filling, the scale also takes into account the level of systolic blood pressure and respiratory rate, and its nature - with or without the involvement of auxiliary muscles in the breathing process [211].

The work was continued and in 1989 a revised scale of injury severity appeared - RTS (Revised Trauma Score). The method is based on the evaluation of 3 parameters: the sum of the scores on the GCS, systolic blood pressure, and respiratory rate. In this scale, there is no need to assess the level of capillary filling or the participation of auxiliary muscles in the breathing process, which are difficult to assess in the field. The authors proposed two variants of the new scale: a simplified sorting variant - T-RTS (Triage Revised Trauma Score) - for use at the prehospital stage and a basic version of RTS - for assessing the outcome of injury and its severity. The triage version is based

on the summation of the scores of the values obtained. The main variant is calculated using the logistic regression equation and is often used to estimate the probability of survival [104].

One of the most common non-specific methods of assessing the severity of the physiological status in clinical practice worldwide, based on the assessment of physiological indicators, are the APACHE II scale (Acute Physiology and Chronic Health Evaluation II) and the simplified scale of acute physiological disorders - SAPS II (simplified acute physiology score II) [163].

There are four variants of the APACHE scale. The most well-known of them was the second version - APACHE II. The second revised version of the APACHE II scale includes 12 parameters: GCS value, heart rate, blood pressure, respiratory rate, rectal temperature, hematocrit, partial pressure of oxygen in arterial blood (PaO₂), white blood cell count, sodium, potassium, plasma creatinine, and arterial pH. Points are also added to the total score for the need for surgical interventions, age, and the presence of chronic diseases [111].

In 1993, a simplified assessment scale of acute physiological disorders (SAPS II) was presented for patients staying in the ICU. The SAPS II scale includes 17 variables: 12 physiological indicators, age, type of hospitalisation (planned surgical, unplanned surgical, or non-surgical) and three categorical variables associated with concomitant diseases (acquired immunodeficiency syndrome, metastatic cancer, and blood cancer) [119].

According to V.V. Aghajanian and others, the use of APACHE and SAPS limits the inclusion of parameters such as blood Na⁺, K⁺ concentration, venous blood plasma bicarbonates, and arterial gas values, as determining these values over time is not available in many health institutions [1, 41].

To assess the severity of multiple organ failure and the likelihood of death in the ICU, a scale of sequential assessment of organ dysfunction - SOFA (Sequential Organ Failure Assessment) is used. The scale was first introduced in 1996 under the name 'Sepsis-related Organ Failure Assessment', but was subsequently renamed after its effectiveness was established in the absence of sepsis. SOFA assesses respiratory

function (arterial partial oxygen pressure (PO₂) to inspiratory oxygen fraction (FiO₂)), coagulation (platelet count), liver function (total bilirubin), cardiovascular function (hypotension), central nervous system (GCS value) and renal function (creatinine or diuresis). Each parameter is evaluated in points, from 1 to 4. Based on the sum of the scores of all indicators, the probability of survival of the patient is determined [207].

Researchers' opinions on the effectiveness of universal and non-specific methods for assessing the severity of the physiological status differ. Some researchers report the effectiveness of methods such as SOFA, APACHE II and SAPS II [93, 128, 172, 176], while others point to their inaccuracy when comparing the results of use for patients with somatic pathologies and for patients with severe trauma [186].

1.2.1.3. Combined Methods for Assessing the Severity of Injuries

Combined methods take into account both the degree of deviation of physiological parameters from the norm and the severity of morphological disorders due to the trauma. In this regard, combined methods, unlike many physiological methods, are characterised by specificity.

Among the first combined methods is that of Yu. N. Tsibin. This method was developed in the late 70s on the basis of Dzhanelidze Research Institute of Emergency Medicine. The method allows to predict the time in hours from the moment of injury to the onset of haemodynamic stabilisation and thereby assess the probability of survival of the patient. To predict the outcome of an injury, the method uses the following parameters: age, pulse rate, systolic blood pressure, and the sum of severity of injuries. The assessment is based on the logistic regression equation [95, 96].

Some authors point out the limitations of this method due to the fact that it does not take into account the severity of the trauma, which is a frequent component of a co-injury [60], the level of consciousness of the patient [52], and such severe injuries as cerebral compression and spinal cord injury, which, in some cases, determine the outcome of the injury [81].

H.J. Oestern et al. in 1985 presented a scale for assessing the severity of polytrauma - PTS (Polytrauma Score, Hannover). The scale assesses the severity of injuries to five anatomical areas of the body and the age of the patient. A modified version of the scale was also presented, which takes into account the sum of points on the GCS, the level of base deficiency, and the Horovitz coefficient: PO_2/FiO_2 . H.J. Oestern et al. suggest classifying polytrauma by severity based on this scale into four degrees: first degree of severity (up to 20 points), second degree (20-34 points), third degree (35-48 points), and fourth degree (>48 points). The predicted mortality rate for each degree is $<10\%$, $<25\%$, $<50\%$, and $<75\%$, respectively [123].

C.R. Boyd et al. proposed a method of assessing the severity of trauma and injury - TRISS (Trauma and Injury Severity Score) to assess the severity of the injury and the physiological status of the patients. The TRISS method is based on two previously created scales for assessing the severity of damage - ISS and RTS. The method also takes into account the influence of the age of the patients (≥ 55 years) on the severity of their physiological status [117].

The second revised classification of the severity of injuries - RISC II (Revised Injury Severity Classification II) for predicting the probability of death in patients with concomitant trauma was proposed by R. Lefering et al. The RISC II scale includes the following parameters: the severity of the two most severe injuries and head injuries on the AIS scale, age, gender, assessment of motor reaction on the scale, pupil reaction and size, the nature of the injury (blunt or penetrating), the sum of points on the scale of the American Society of Anesthesiologists - ASA, the need for cardiopulmonary resuscitation, systolic blood pressure, INR, and deficiency of blood bases and haemoglobin [214].

Separate inclusion of the severity of head injuries and the severity of injuries to other anatomical areas in the RISC II prognostic model contributed to its prognostic power [214].

A Severity Characterisation of Trauma (ASCOT) scale was introduced in 1990. To describe the severity of damage, the scale uses the codes described in the AIS scale (updated in 1985). The scale also includes the following physiological parameters: the

level of consciousness according to the GCS upon admission to the ICU, the patient's age, systolic blood pressure, and respiratory rate [103].

Among domestic methods of assessing injury severity, the military field surgery (MFS) method of objective assessment of injury severity became widely used. This method was developed in 1992 and includes several scales, based on both anatomical and physiological principles and adapted to meet specific clinical needs. The method includes: scales for assessment of the severity of injuries in three versions: for gunshot wounds, for non-gunshot wounds, and for mechanical injuries; the MFS triage scale; scales for assessment of the physiological status of the injured or wounded person in three versions: on admission, for dynamic follow-up in medium-level institutions, and in specialised centres; scales for prediction and diagnosis of fat embolism; a scale for diagnosis of cardiac contusion; a surgical tactics scale; and an endovideosurgical scale. The authors also presented qualitative equivalents of scores on scales; the predicted frequency of complications; the probability of death and the duration of disability [20, 26].

The MFS Injury Scale (mechanical injury) includes 84 of the most common injuries in clinical practice in six anatomical areas (head, chest, spine, abdomen, pelvis, and limbs) with severity codes from 0.05 to 19. The severity of the injuries includes the sum of all existing injuries in all anatomical areas, which undoubtedly increases the accuracy of assessment in multiple and concomitant injuries [26].

The MFS Scale - admission status scale is used to assess the severity of the physiological status of the patient upon admission to hospital. The scale includes 12 physiological parameters: the colour of the skin, the nature of external respiration, auscultative changes in the lungs, reaction to pain, speech contact, systolic blood pressure, pulse rate, pulse pattern, pupillary or corneal reflexes, pupil size, approximate amount of blood loss, and intestinal peristalsis noises [20].

According to A. N. Tulupova et al., the VPH method was developed on the basis of a clinical material of a sample population, the main component of which were healthy persons (service men) of a limited age category, and therefore this method does

not always give an accurate assessment of the severity of injuries in persons of other age groups suffering from concomitant diseases [69].

According to A.V. Semenova et al., the following points can be attributed to the disadvantage of the method of assessing the severity of VPH injuries: the sign used in different scales is encoded with different values, which can lead to evaluation errors; the level of consciousness of the patient is not evaluated upon admission; a large number of parameters used makes the process of assessing the severity of injuries difficult and takes a lot of time; the use of subjective parameters, such as the noise of intestinal peristalsis, the approximate amount of blood loss, and the nature of external respiration [81].

1.2.2. Analysis of the Validity of Methods for Assessing Severity and Prognosis of Immediate Trauma Outcomes

Judging by the number of scientific publications, the methods described above are the most common in clinical practice and the most discussed in the scientific field. But the question of which of the methods is the most effective and accurate is a very controversial issue. The analysis of special literature indicates the presence of many domestic and foreign publications devoted to the study of the effectiveness of various methods [22, 75, 101, 108, 125, 126, 127, 171, 191, 199]. Below we will present a small part of the published scientific material on the study of this problem.

The most common parameter used in evaluating the effectiveness of severity assessment methods is the accuracy of predicting the probability of a lethal outcome. Most researchers conduct an assessment by constructing a ROC curve (Receiver Operating Characteristic), followed by measuring the area under the curve and evaluating the sensitivity and specificity of the method. The predicted and actual deaths of the patients are also compared. Table 1 presents the results of comparative studies of the prognostic value of various methods of assessing severity.

Table 1. The results of scientific research on the comparative analysis of the accuracy of some methods of assessing the severity of predicting the fatal outcome of injuries

Studies	Compared methods	Identified results
[157, 161, 192]	NISS and ISS	NISS is superior to ISS
[206]	APACHE II, NISS, and ISS	APACHE II is superior to NISS and ISS
[129]	APACHE II and GCS	APACHE II is superior to GCS
[130]	APACHE II and SOFA	Scales equal in accuracy
[67]	APACHE II and ISS	ISS is superior to APACHE II
[68]	APACHE II and ISS	Scales equal in accuracy
[176]	SAPS II and SOFA	SAPS II is superior to SOFA
[130, 152, 195, 212]	TRISS APACHE II and SOFA	Scales equal in accuracy
[102, 154]	TMPM, ISS, NISS, AIS, and ICISS	TMPM is superior to the rest
[149]	ASCOT and TRISS	ASCOT surpasses TRISS
[208]	ASCOT and TRISS	Scales equal in accuracy
[9, 67]	ISS, APACHE II, MFS-I(injury), and MFS-CA (physiological status on admission)	ISS and APACHE II are superior to MFS-I and MFS-CA
[166]	RISC II and TRISS	RISC II is superior to TRISS
[101]	ISS, APS, NISS, and ICISS-9	Scales equal in accuracy
[148]	ICISS-9 and ISS	ICISS-9 is superior to ISS
[199]	ICISS-9, ISS, and TRISS	ICISS-9 is superior to ISS and TRISS
[126]	APACHE II and TRISS	APACHE II is inferior to TRISS
[108]	APACHE II and SAPS II	Scales equal in accuracy
[33, 59]	ISS and MFS-I (MT - mechanical trauma)	Scales equal in accuracy
[155]	APACHE III and SAPS II	Scales equal in accuracy
[22]	ASCOT and SAPS II	Scales not suitable for this purpose

The data in Table 1 show that the published research results are extremely contradictory. The superiority of one method, revealed in one study, is rejected in another. When analysing scientific publications comparing the ICISS, NISS, and ISS

scales on the accuracy of predicting the fatal outcome in patients with combined trauma, H. Tohira et al. obtained extremely contradictory results [192].

The clinical material used in the development of non-specific methods includes both trauma patients and patients with general somatic diseases. For example, about 30% of the clinical material used in developing the APACHE II scale relates to trauma patients, and the bulk of the clinical material relates to patients with somatic diseases [111].

Some authors note that the use of a combination of trauma-specific and non-specific methods for assessing the severity of the physiological status can increase the accuracy of predicting the fatal outcome of patients [176], but non-specific methods do not take into account differences in the course of decompensated forms of somatic diseases and traumatic disease.

S.K. Park et al. when studying the effectiveness of the APACHE II and SAPS II scales in predicting mortality in patients with severe TBI revealed that the mortality predicted by both methods (APACHE II= 37.7%; SAPS II= 38.4%) is almost 12% higher than the actual mortality (24.8%). The authors identify the need for a new method of predicting the outcome of a traumatic brain injury, taking into account the severity of brain dysfunction [108].

Some researchers point to differences in the course of trauma illness in concomitant injuries depending on the localisation of the leading injury and to a decrease in the accuracy of common trauma-specific methods, such as RISC II, in patients with severe traumatic brain injury [216].

Thus, according to the literature review, a large number of specific and non-specific methods have been proposed to assess the severity and predict outcomes of injuries. Nevertheless, the results of the studies during testing and comparison turned out to be ambiguous and often mutually exclusive. Researchers have often observed a decrease in the accuracy of these methods and scales in patients with severe TBI, and the problem of predicting the immediate outcomes of combined midface trauma based on specific clinical material has not been studied.

1.3. Current Status of the Problem of Treating Survivors with Concomitant Midface Trauma

The analysis of special scientific literature available to us shows that most of the studies devoted to the topic of midface injury were aimed at studying the functional and aesthetic consequences of isolated injuries. The problem of concomitant midface trauma and craniofacial trauma as a whole has received much less attention.

A significant part of the work on optimising treatment tactics in patients with concomitant midface trauma was carried out on the basis of the departments of Maxillofacial Surgery and Surgical Dentistry of Kirov Military Medical Academy and St Petersburg University. It included the development and testing of the concept of multi-stage surgical treatment (MST) and the development of devices for external fixation of fractures of the maxillofacial bones.

I.M. Belous studied the possibility of using the MFS method of severity assessment in the treatment of patients with a concomitant maxillofacial trauma. The author also presented a scale for assessing the severity of maxillofacial injury 'MFS - MFI (MT - mechanical injury) based on the MFS method. According to the author, the periods of fixation of fractures are determined depending on the degree of compensation for the physiological status of the patient. The author noted the positive results of applying the MFS method in the organisation of treatment tactics for patients, such as a 2.37% reduction in mortality, a 2-fold reduction in complication rates, and a 2-fold reduction in length of stay in the ICU and in hospital in general) [10].

M.O. Danilevich proposed to use the method of the revised trauma score - RTS to assess the severity of craniofacial trauma. The author points out the simplicity and convenience of using this method, as well as its superiority in accuracy compared to the GCS. The accuracy of prediction of the RTS method, according to the author, is 85% [28].

According to D.Yu. Madai et al., the use of tactics based on the MFS method of objective assessment of trauma severity and the concept of traumatic illness has reduced mortality in patients with concomitant craniofacial trauma by 5.4 times, the

number of complications in this group of patients - by 1.4 times, and the length of stay in the ICU - by 1.6 times [58, 61, 64, 89].

K.P. Golovko et al. established that diagnostics and surgical tactics based on the application of the MFS method of objective assessment of injury severity, endovideosurgery, and minimally invasive extrafacial osteosynthesis in treatment of patients with combined traumas of maxillofacial region, have contributed to the 61% decrease of rates of purulent-septic complications, 37.5% decrease of lethality, and 1.8 times (from 6.9 ± 0.4 to 4.1 ± 0.7 days) decrease of ICU-treatment duration [24, 56].

From the literature review data, it can be seen that the amount of work done to optimise treatment tactics for survivors of concomitant midface trauma based on objective assessment of injury severity is limited. Most researchers use the MFS injury severity assessment method to assess the severity of injuries. In addition, in previous studies, insufficient attention was paid to the problem of treating patients with extremely severe concomitant midface trauma.

CHAPTER 2. MATERIALS AND METHODS OF RESEARCH

2.1. Research material

The study was conducted at the clinical facility of the Department of Maxillofacial Surgery and Surgical Dentistry of St Petersburg University - the 1st Neurosurgical Department (for patients with concomitant craniofacial trauma and hearing and vision injuries), and the Intensive Care Unit No 2 and the Concomitant Trauma Department of St Petersburg Aleksandrovskaya Hospital. The study is based on the data of clinical material presented by 111 patients with concomitant midface trauma. The criteria for inclusion in the study were: concomitant midface injury, severity of injuries ≥ 1 point on the MFS - I (MT) scale (severe or extremely severe), duration of hospital treatment $>$ of 3 days (patients who died in the first 3 days after the injury were excluded from the study).

The proportion of male patients is 74.8% (n=83) of all patients; the proportion of female patients is 25.2% (n=28). The age of the patients varies from 18 to 96 years. The main proportion of patients is represented by young people (18-44 years old) - 55% (n=61). Middle-aged patients (45-59 years old) comprised 19.8% (n=22) of the total number; elderly patients (60-74 years old) - 16.2% (n=18); old patients (75-89 years old) - 8.1% (n=9); long-lived patients (over 90 years old) - 0.9% (n=1). The average age of all the patients was 46.9 ± 18.5 years. The age distribution of the patients is shown in detail in Figure 1.




Figure1. Distribution of the studied patients by age.

According to the conclusion of the staff therapist, in the group of 95 patients, 52 patients (54.7%) suffered from concomitant diseases. Diseases of the cardiovascular system were diagnosed in 44 (46.3%) patients; nervous system diseases in 20 (21.1%) patients; urinary system diseases - in 8 (8.4%); endocrine system diseases - in 7 (7.4%); digestive system diseases - in 5 (5.3%); and respiratory diseases - in 4 (4.2%) patients.

Assaults with the use of physical force were the most common cause of injuries and occurred in 38 (34.2%) patients. In second place is the car accident, with 25 (22.5%) patients. Fall-related injury was observed in 24 (21.6%) patients. 14 (12.6%) of the patients fell from heights over 3 metres and 10 (9%) fell from their own height. Two (1.8%) patients were injured with traumatic weapons. The mechanism of injury could not be established in 22 (19.8%) patients. The mechanisms of injury occurrence are presented in Table 2.

Table 2. Mechanisms of occurrence of concomitant midface injury

Injury mechanism	Number of patients, n = 111	
	absolute number	%
Assault with the use of physical force	38	34.2
Traffic accident	25	22.5
Falling from a height of over 3 metres	14	12.6
Falling from one's own height	10	9
Injury from a traumatic weapon	2	1.8
Injury under unknown circumstances	22	19.8

In addition to midface injuries, all the patients (n=111) were diagnosed with TBI. Open TBI was diagnosed in 70 (63,1%) patients and closed TBI - in 41 (36,9%). Injuries in the chest area occurred in 37 (33.3%) patients; limb injuries - in 33 (29.7%); abdomen - in 10 (9%); spine - in 9 (8.1%); pelvis - in 6 (5.4%); and neck - in 3 (2.7%) patients. On average, without taking into account the midface injury, each patient was diagnosed with damage to 1.9 anatomical areas. The distribution of injuries by anatomical areas in the examined patients is presented in Table 3.

Table 3. Distribution of injuries by anatomical areas in patients with concomitant midface trauma

Anatomical area	Number of patients, n=111	
	absolute number	%
Head (TBI)	111	100
Neck	3	2.7
Chest	37	33.3
Stomach	10	9
Pelvis	6	5.4
Spine	9	8.1
Limbs	33	29.7

Injuries in the head area were assessed as injuries leading in severity in 91.9% of cases (in 102 patients); chest injuries, as well as limb injuries - in 2.7% (in 3 patients,


respectively); spine area - in 1.8% (in 2 patients); abdominal area - in 0.9% (the 1 patient).

Among the fractures of the midface bones, the first place is occupied by fractures of the eye socket walls, which were diagnosed in 61 (55%) patients. The second place is occupied by fractures of the upper jaw - in 43 (38.7%) patients. Fractures of the upper jaw of the upper type were observed in 25 (22.5%) patients; of the middle type - in 29 (26.1%); of the lower type - in 18 (16.2%). Fractures of the zygomatic bone occurred in 40 (36%) patients; of nasal bones - in 37 (33.3%) patients. Soft tissue injuries of the midface region were diagnosed in 66 (59.5%) patients. The distribution of midface injuries in the examined patients is presented in Table 4.


Table 4. Distribution of midface injuries in the examined patients

Anatomical location	Number of patients, n=111	
	absolute number	%
Upper jaw	43	38.7
Eye socket	61	55
Zygomatic bone	40	36
Nose bones	37	33.3
Soft tissues	66	59.5

The severity of injuries on the MFS-I 'MT' scale in the examined patients ranged from 1 to 24.1 points (median = 12.2 points); on the ISS scale - from 9 to 50 points (median =29 points). 54 survivors had severe injuries (1 to 12 on the MFS-I (MT) scale); 57 survivors had extremely severe injuries (≥ 12 points on the MFS-I (MT) scale). The severity of injuries in the examined patients is shown in Figure 2.



a



b

Figure 2. Severity of injuries in the examined patients: a) on the MFS-I (MT) scale; b) on the ISS scale.

In the group of 95 patients, traumatic shock upon admission to the clinic was diagnosed in 78.9% (n=75) of cases. Traumatic shock of the second degree was observed in 36 of 75 patients (48%); of the first degree in 30 (40%); and of the third

degree in 9 (12%). Table 5 shows the distribution of patients by severity of shock, indicating the approximate amount of blood loss and the value of the Allgöwer shock index.

Table 5. Distribution of patients by severity of shock

Severity of shock	Number of patients, (n=75)		Approximate blood loss, l	Allgöwer shock index (M±SD)
	absolute number	%		
I degree	30	40	0,5-1	0,9±0,3
II degree	36	48	1-1,5	1,1±0,4
III degree	9	12	More than 1.5	1,3±0,3

2.2 Research methods

2.2.1. Methodology of Data Collection

When collecting clinical material, generating a preliminary dataset, and mathematical processing of research data, the author was guided by the general principles of the biometric analysis of medical data described by Professor D.Yu. Madai and Candidate of Physics and Mathematics A.G. Barth et al. in the textbook 'Biometric Analysis of the Effectiveness of Treatment' [12].

The methodology involves measuring a feature or features under study at several predetermined and fixed time points for all the patients involved in the study. These time points in our study corresponded to the 1st, 3rd, 7th, 14th days after the injury. These time points were chosen taking into account the theory of traumatic illness generally accepted in traumatology and correspond to its periods. This technique allows one to observe the dynamics of changes in the studied feature in different periods of traumatic illness, which in turn allows one to study the features and nature of its course. The values of the studied features were also collected for the last day of life of patients with a fatal outcome of combined trauma.

The generated dataset includes 77 qualitative variables and 114 quantitative variables and consists of several data units:

1. The first unit includes general data about the patients, such as data identifying the patients, the age of the patients, the duration of hospital stay, concomitant diseases, and other general data.
2. The second unit contains data on the injury itself, the mechanism of its occurrence, diagnosis, treatment, complications, outcome, and other qualitative data.
3. The third unit is the most massive and includes indicators of clinical and laboratory indicators measured at the above-mentioned five time points.
4. The fourth unit contains data on the severity of injuries and the physiological status of the examined patients.

2.2.2. Radiation methods of examination of patients

The method of multi spiral computed tomography (MSCT) was used as the main method of radiation diagnosis of craniofacial injuries. Computed tomography was performed on all the patients upon admission to hospital. Repeated computed tomography was also performed, if necessary, after the initial stabilisation of the physiological status of the patients and after the final fixation of midface bone fractures.

During the computed tomography, the clinic's standard equipment was used - the Aquilion 16 CT scanner from Toshiba (Japan) with a 16-detector multi-slice scanning system (Registration certificate No. FSZ 2007/00892) and the Somatom Definition AS spiral CT scanner from Siemens (Germany) with a 128-detector multi-slice scanning system (Registration certificate No. FSZ 2008/02797).

2.2.3. Study of Clinical and Laboratory Signs Reflecting the Severity of the Physiological status of the Patients

Measurement of clinical and laboratory physiological signs was carried out using standard equipment of the clinic:

1. The PICCO Plus Haemodynamic Monitor by PULSE (Germany) (Registration certificate No. FS 2005/511): was used to measure heart rate (HR) and blood pressure (BP).

2. Haematological analyser KX -21N by Sysmex (Japan) (Registration certificate No. FSZ 2011/11182) was used to measure the number of leukocytes, lymphocytes, monocytes, and haemoglobin levels in venous blood.

3. Biochemical analysers iLab Taurus by Instrumentation Laboratory (Italy) (Registration certificate No. FSZ 2012/12577), Cobas Integra 400 plus by Roche Diagnostics (Switzerland) (Registration certificate No. FSZ 2012/11531), and Architect c8000 Processing Module by Abbott Laboratories Diagnostics Division and Toshiba Medical Systems Corporation (USA, Japan) (Registration certificate No. REN 2014/2010) were used to measure the level of alanine aminotransferase (ALT), total bilirubin, glucose, urea, as well as concentrations of potassium (K⁺), sodium (Na⁺), and chlorine (Cl⁻) ions in blood.

4. Coagulation analyser CS 5100 e by Sysmex (Japan) (Registration certificate No. RZN 2013/762) was used to determine the international normalised ratio (INR).

5. The analyser of acid-base and gas composition of blood of the ABL800 Flex series by Radiometer Medical ApS (Denmark) (Registration certificate No. RZN 2015/2415) was used to measure the partial pressure of carbon dioxide (pCO₂), partial pressure of oxygen (pO₂), hydrogen index (acidity- pH), the level of oxygen saturation of haemoglobin in arterial blood (sO₂), and blood bicarbonate (cHCO₃) concentrations.

2.2.4. Methods for Assessing the Severity of Concomitant Midface Trauma

Assessment of the severity of injuries in the patients studied was carried out using two widely used methods: the MFS - I (MT) method of injury severity assessment and the ISS injury severity scale.

2.2.4.1. Injury Severity Score MFS-I (MT)

To assess the severity of injuries on this scale, the points for each injury diagnosed in a patient are summed up. The scale contains 84 names for the most common injuries in clinical practice. Table 6 shows the names of injuries in the head area and the number of points assigned to them, according to this scale.

Table 6. Assessment of the severity of injuries on the MFS - I (MT) Head scale [cited by: 20, p. 722]

Item no.	Type of injury	MFS - I
1	Wounds of the soft tissues of the head	0.05
2	Closed fractures of the bones of the nose	0.2
3	Concussion of the brain	0.2
4	Fractures of the jaws	0.3
5	Mild brain contusion	0.3
6	Moderate brain contusion with fractures of the vault of the skull	0.5
7	Moderate brain contusion with closed fractures of the vault and base of the skull	0.6
8	Moderate brain contusion with open fractures of the vault and base of the skull	2
9	Compression of the brain accompanied by mild contusions	7
10	Severe brain contusion with damage to the upper brainstem	12
11	Compression of the brain accompanied by severe contusions	18
12	Severe brain contusion with damage to the lower brainstem	19

The calculated sum of points is subsequently subject to interpretation in the traditional qualitative gradations of severity. Light injuries have quantitative values in the range of 0.05 - 0.49 points; moderate injuries in the range of 0.5 - 0.99 points; and severe injuries in the range of 1.0 - 12.0 points. Injuries with a severity of > 12.0 points are extremely severe (Table 7).

Table 7. Gradation of severity of injuries in patients [cited by: 20, p. 729]

Traditional injury gradation	Quantitative assessment of injuries (score)
Mild	0,05 - 0,49
Moderate	0,5 - 0,99
Severe	1,0 - 12,0
Extremely severe	> 12,0

2.2.4.2. Injury Severity Score ISS

To calculate the severity of injuries on the ISS scale, the scoring method proposed in the AIS scale is used. According to the AIS scale, each injury, depending on its severity, is assigned a certain number of points. Mild injury is estimated at 1 point; moderate injury at 2 points; severe injury without a threat to life at 3 points; severe injury with a threat to life at 4 points; critical injury at 5 points; fatal injury (injury incompatible with life) at 6 points (Table 8) [202].

Table 8. Qualitative and quantitative gradation of injury severity on the AIS scale

Gradation of injury severity	Value in points
Mild injury	1
Moderate injury	2
Severe injury without threat to life	3
Severe life-threatening injury	4
Critical injury	5
Fatal injury - incompatible with life	6

To calculate the severity of injuries on the ISS scale, add up the squares of points on the AIS scale of the three most severe injuries in three different anatomical areas. According to the AIS and ISS scales, injuries are assessed in six anatomical areas: head and neck, face, chest, limbs and pelvic organs, abdomen, and external integuments [202].

Thus, the sum of points on the ISS scale varies from 1 to 75. In the presence of one fatal injury incompatible with life, 75 points are automatically assigned to it, despite the severity of other injuries [202].

There are no clear quantitative boundaries in the ISS scale, which are necessary for interpreting the sum of points obtained in assessing severity into qualitative gradations of severity. However, in the field of traumatology, it is customary to attribute injuries with a score of 17 >to severe and polytrauma [25].

2.2.5. Statistical Processing of Research Data

Statistical processing of the research data was carried out using the STATISTICA 10.0 applied software package together with Associate Professor of the Department of General Mathematics and Computer Science of St Petersburg University V.A. Bart. When describing the results of the study, the author was guided by the recommendations and requirements of leading modern specialists in presenting the results of medical research and, in particular, the recommendations of Professor T.A. Lang [3, 42].

At the stage of exploratory analysis of the study data, it was found that the vast majority of the studied quantitative features had a normal or logarithmically normal distribution. Measures of the central trend and the range of features with a normal distribution will be presented in the format of the mean (M) and standard deviation (SD); features with a logarithmic normal distribution - in the form of median (Mdn), minimum (Min), and maximum values (Max). In order to unify the format for presenting descriptive statistics of features, measures of central trend and scope in the section 'exploration analysis' will be presented in the form of median (Mdn), minimum

(Min), and maximum (Max) values. The hypothesis of the equality of means across groups was tested using the Mann-Whitney test or the Student's t-test.

Factor analysis (the principal component method) was used as the main method of identifying syndrome complexes (factors) characterising the severity of the physiological status of patients in the last day before the onset of death, as well as to determine the optimal time point for predicting the nearest outcomes. The use of factor analysis made it possible to reduce the total number of signs without significant loss of information.

Correlation analysis and correlation pleiades were used to assess the strength and significance of the interrelationships of the features forming the selected factors.

Discriminant analysis was applied in the creation of a mathematical model for predicting the immediate outcomes of concomitant midface trauma.

CHAPTER 3.

EXPLORATORY ANALYSIS OF RESEARCH PAPER DATA

3.1. Exploratory Analysis of the Research Material

To address the research objectives, three subsets were formed from the total dataset (n=111 patients), which will be referred to as the first, second, and third subsets of data, respectively:

1. The first data subset. 95 patients were included in this data subset. 22 physiological signs reflecting the severity of their physiological status in the course of traumatic disease were studied in the patients of this subset. The patients were divided into 2 groups depending on the immediate outcomes of the concomitant midface injury. The first group – patients with a favourable outcome (n=50 (52.6%)); the second group – patients with an unfavourable outcome (n=45 (47.4%)). We used the data of the first subset to address the first and second tasks of this study (searching for syndrome complexes and signs characterising the severity of the physiological status of patients with concomitant midface trauma in the course of traumatic disease; developing a methodology for predicting the immediate outcomes of concomitant midface trauma).

2. The second data subset. It includes 48 patients with severe concomitant midface trauma (from 1 to 12 points on the MFS - I (MT) scale). Based on the data of the second subset, two groups were created: the first group (conventionally called the first group of observed patients) – includes 27 (56.25%) patients, whose treatment was carried out using the MST tactics, taking into account the severity of their physiological status; the second group (conventionally called the first retrospective group) – includes 21 (43.75%) patients, whose treatment was carried out by the traditional clinical method.

3. The third data subset. It includes 57 patients with extremely severe concomitant midface trauma (≥ 12 points on the MFS - I (MT) scale). The patients of the third data set were divided into two groups: the first group (conventionally called the second group of observed patients) – includes 21 (36.8%) patients, whose treatment

was carried out, as in the patients of the first group of observed patients, using the MST tactics, taking into account the overall severity of their physiological status; the second group (conventionally called the second retrospective group) – includes 36 (63.2%) patients treated with the traditional clinical method. We used data from the second and third subsets to address the third task of the study (evaluation of the effectiveness of the MST tactic in patients with severe and extremely severe concomitant midface trauma).

The formation of two test subsets from the dataset was carried out to address the third research objective, aiming to maximize the preservation of homogeneity within the comparison groups based on the severity of the trauma. Figure 3 shows how the number of patients of the three data subsets has evolved over time.




Figure 3. Change in the number of patients of the three data sets over time.

3.2. Exploratory Analysis of Studied Physiological Signs

The severity of the physiological status of patients can be measured by the degree of deviation of physiological signs from the physiological norm. The greater the magnitude of these deviations, the more severe the physiological status.

Observation time is an important component of any medical research. B.B. Bondarenko et al. described four variants of the time of change in the sign values from the observer's point of view: 1. time of examination; 2. calendar time; 3. time of the sign; 4. time of the course of the disease (most often used) [4]. In our study, the signs were measured in two time variants. The first variant is the time of the course of the traumatic illness (1st, 3rd, 7th, and 14th days from the moment of injury), and the second variant is the time of waiting for the outcome (on the last day before the fatal outcome (only in the group of patients with an unfavourable outcome). The time point in the second variant of the observation time will be designated as day -1.

Below we provide a list of the signs we are investigating (22 signs) with the indication of the number and code of each one. A brief description of the physiological significance of the signs is also given.

BT / (1) The body temperature of the patients (36.8°C) - hyperthermia may indicate general intoxication as a result of the development of local, visceral, or generalised infectious complications. Spontaneous hypothermia often occurs on the first day after injury, especially if the injury is accompanied by massive blood loss.

HR / (2) Heart rate (from 60 to 100 beats per minute) characterises cardiac activity and the state of haemodynamics. Tachycardia is a sign of general intoxication with the development of infectious complications in patients.

SBP/ (3) Systolic blood pressure (120 mmHg) reflects the state of haemodynamics and the state of the cardiovascular system. A decrease in systolic blood pressure to 80 mmHg and below indicates the instability of haemodynamics and is an indication for inotropic support. If, at the time of measuring the sign, inotropic support was provided to the patient, then systolic blood pressure was automatically assessed at 80 mmHg.

Hb / (4) Haemoglobin (120-140 g / l for women; 130-160 g / l for men) - acute posthemorrhagic anemia occurs as a direct result of mechanical damage to tissues and organs, accompanied by massive blood loss. The process of erythropoiesis and the level of haemoglobin can also be affected by the development of infectious and purulent-

septic complications in patients. Haemoglobin is also an indirect sign of oxygenation of tissues and organs.

WBC / (5) Leukocytes ($4-8.8 \times 10^9 / l$) - the level of leukocytes reflects the state of immune-reactive processes and resistance of the body during the development of infectious complications. Interpretation of the values of this sign should be carried out taking into account other signs of infectious complications, since patients with severe trauma often develop redistributive leukocytosis, especially in the first and second periods of traumatic illness.

Lym/ (6) Lymphocytes ($1,2-3,0 \times 10^9 / l$) are the main cells of the immune system responsible for humoral and cellular immunity. Reflect the state of resistance of the body and the nature of the immune response in the development of infectious complications.

Mono/ (7) Monocytes ($0.29-0.6 \times 10^9 / l$) are cells of the immune system involved in the process of phagocytosis in infectious complications.

NLR/ (8) Neutrophil-Lymphocyte Ratio (0.78-3.5) is an indicator of the morphological composition of leukocytes, reflecting the severity of the 'stress load' of the immune system [220]. It is calculated by dividing the absolute number of neutrophils by the number of lymphocytes.

$$NLR = \frac{\text{number of neutrophils}}{\text{number of lymphocyte}}, \quad (1)$$

The greater the 'stress load' of the immune system, the greater the number of neutrophils, the smaller the number of lymphocytes and the higher the NLR index [221].

LII/ (9) Leukocyte Intoxication Index (LII) according to Ya.Ya. Kalf-Kalif (0.5-1.0) - reflects the severity of the infectious process and endogenous intoxication. An increase in LII to 4.0 or more is considered a sign of significant endogenous intoxication [39]. LII is calculated by the following formula:

$$LII = \frac{(S+2R+3Y+4MYC)(PC+1)}{(Mono+Lym)(Eo+1)}, \quad (2)$$

where S – segmented neutrophils;

R – rod-shaped neutrophils;

Y – young;

MYC – myelocytes;

PC – plasma cells;

Mono – monocytes;

Lym – lymphocytes;

Eo – eosinophils.

ALT / (10) Alanine aminotransferase (≤ 34 U/l for women; ≤ 45 U/l for men) is an indicator of the severity of damage to liver cells in violation of its function and the development of organ failure.

Bil./ (11) Total blood bilirubin (2.5- 20.5 mmol/L) - reflects the functional state of the liver. Hyperbilirubinemia in patients with severe trauma may be the result of increased haemolysis in infectious and purulent-septic complications or impaired liver function and organ failure.

BG / (12) Blood glucose (3.89- 6 mmol /L) - stress hyperglycemia often develops with a complicated course and an unfavourable outcome of critical physiological status. The mechanism of its development is associated with hypersecretion of glucocorticoids and catecholamines, which lead to activation of lipolysis, gluconeogenesis, inhibition of aerobic glycolysis, suppression of insulin secretion, and the development of insulin resistance [37].

Na+ / (13) Concentration of sodium ions in the blood (136-145 mmol/L) - sodium ions are the main cation of extracellular fluid, which plays an important role in regulating the water balance, maintaining blood pressure, conducting nerve impulses, and muscle contraction. Hypernatremia can occur against the background of hypovolemia as a result of blood loss and a decrease in the volume of circulating blood. Hypernatremia can also be iatrogenic as a result of excessive administration of sodium

ions, a tactic that is sometimes used to combat cerebral oedema [144]. Dysnatremia is an indirect sign of impaired function of the excretory system and the development of renal failure.

K^{+} / (14) Concentration of potassium ions in the blood (3.5- 5.1 mmol/L) - potassium ions are the main cation of the intracellular fluid whose main function is the regulation of neuromuscular excitability. Violation of the concentration of potassium ions in the body has a negative effect on the work of the respiratory muscles and cardiac activity. The level of potassium in the blood is regulated mainly by renal excretion under the action of the hormone aldosterone, therefore, a violation of potassium concentration may indicate kidney dysfunction and the development of renal failure.

Cl^{-} / (15) The concentration of chlorine ions in the blood (98- 107 mmol/L) is the main anion of extracellular fluid, participates in the regulation of osmotic pressure, muscle activity, and acid-base balance. An indirect sign of a violation of the function of the excretory system and the development of metabolic acidosis.

BU / (16) Urea in the blood (3- 9.2 mmol/L) is the final product of protein breakdown and an indicator of the functional state of the excretory system.

INR/ (17) International normalised ratio (0.8- 1.2) is a standardised indicator of blood clotting. Reflects the severity of post-traumatic coagulopathy [6, 18].

pH/ (18): hydrogen index (7,350-7,450) is an indicator of blood acidity and acid-base balance. Blood buffer systems, lungs, and kidneys are involved in the regulation of pH levels. Deviation of pH from the physiological norm indicates the development of acidosis or alkalosis. Both physiological statuses may have respiratory (with reduced respiratory efficiency) or metabolic (with impaired renal function) mechanisms of development. To differentiate the aetiology of acidosis or alkalosis, the pH level must be evaluated together with the partial pressure of carbon dioxide (pCO_2) and the level of blood bicarbonate (HCO_3^{-}).

pCO_2 / (19): partial pressure of carbon dioxide in arterial blood (32-48 mmHg) is an indicator characterising the adequacy of pulmonary ventilation and the respiratory link of the acid-base state. The pCO_2 level is used to determine the type of respiratory failure and to establish the nature and aetiology of acid-base disorders.

pO₂/ (20): partial pressure of oxygen in arterial blood (83 - 108 mmHg) - reflects the adequacy of the process of oxygenation of blood in the lungs, and the amount of oxygen available to the body, not related to haemoglobin. pO₂ is used in the diagnosis of hypoxemia and respiratory failure.

SO₂/ (21): Haemoglobin oxygen saturation in arterial blood (95-99%) is an important sign of blood oxygenation. Oxygen associated with haemoglobin accounts for up to 98% of the total amount of oxygen in the blood. pO₂ is a glycemic factor that determines the level of SO₂. The level of haemoglobin oxygen saturation is used in the diagnosis of respiratory failure.

HCO₃⁻/ (22): Blood bicarbonate level (21.2-27 mmol/L for women; 22.2-28.3 mmol/L for men) is a metabolic component of the acid-base state and the main component of the blood buffer system. The HCO₃ level is used in the diagnosis of the nature and aetiology of acid-base balance disorders. Violation of HCO₃ concentration is also an indirect sign of renal failure.

Table 9 provides information on the number of studies conducted on each sign at each time point. It should be pointed out that the number of patients examined at each time point and the number of studies conducted on each attribute may differ. This is due to organisational, technical, and financial reasons, due, among other things, to the policy of managing patients with severe trauma at the clinical facility of the study. For the same reasons, signs relating to blood gas analysis, which is only carried out in the ICU wards, have been investigated much less frequently than others.

Table 9. The number of sign measurements performed at each time point

Sign		Number of studies at each time point				
No.	Code	Day 1	Day 3	Day 7	Day 14	Day -1
1	BT	95	95	89	57	41
2	HR	95	95	89	57	44
3	SBP	95	95	89	57	44
4	Hb	92	74	51	28	36
5	WBC	91	73	51	28	36
6	Lym	86	73	51	28	36
7	Mono	86	73	51	28	36

Continuation of Table 9

8	NLR	86	73	51	28	36
9	LII	86	73	51	28	36
10	ALT	82	70	52	30	36
11	B	79	60	44	24	36
12	BG	90	72	52	28	35
13	Na ⁺	80	68	51	24	36
14	K ⁺	80	68	51	24	36
15	Cl ⁻	79	68	50	24	36
16	BU	79	71	51	29	36
17	INR	79	65	49	26	34
18	pH	42	41	31	14	23
19	pCO ₂	42	41	31	14	23
20	pO ₂	42	41	31	14	23
21	SO ₂	42	41	31	14	23
22	HCO ₃ ⁻	42	41	31	14	23

Tables 10, 11, 12, 13, and 14 present descriptive statistics of the studied signs in patients with favourable and unfavourable outcomes. The p-level of significance of differences in the mean values of the studied signs by groups using the Mann-Whitney test is also given (taking into account Bonferroni's corrections for the multiplicity of comparisons).

Table 10. Descriptive statistics of signs on Day 1

Sign		Favourable		Unfavourable		Mann-Whitney Test, P-level
No.	Code	n	Median [Min.; Max.]	n	Median [Min.; Max.]	
1	BT	50	36,80 [36,00; 37,70]	45	36,70 [35,90; 38,60]	0.979
2	HR	50	101,00 [70,00; 160,00]	45	110,00 [62,00; 151,00]	0.200
3	SBP	50	113,00 [60,00; 160,00]	45	111,00 [70,00; 160,00]	0.841
4	Hb	49	117,00 [54,00; 154,00]	43	104,00 [60,00; 153,00]	0.107
5	WBC	48	16,35 [4,87; 27,27]	43	13,56 [4,88; 24,06]	0.040
6	Lym	44	1,44 [0,19; 3,31]	42	0,67 [0,11; 2,75]	0.001
7	Mono	44	1,00 [0,09; 2,33]	42	0,74 [0,10; 2,08]	0.141
8	NLR	44	9,17 [0,96; 31,33]	42	17,05 [0,98; 46,94]	0.012
9	LII	44	5,60 [1,05; 13,08]	42	7,05 [0,87; 19,98]	0.030
10	ALT	37	47,00 [17,00; 167,00]	45	33,00 [8,00; 116,00]	0.042

Continuation of Table 10

11	B	35	14,20 [1,90; 33,80]	44	12,85 [6,00; 31,20]	0.805
12	BG	47	7,85 [4,61; 14,10]	43	9,59 [5,02; 15,75]	0.004
13	Na+	36	139,00 [132,00; 149,00]	44	141,00 [133,00; 155,00]	0.227
14	K+	36	3,80 [2,60; 5,70]	44	3,80 [2,52; 5,10]	0.568
15	Cl-	36	108,50 [99,00; 117,00]	43	106,00 [96,00; 116,00]	0.386
16	BU	36	4,00 [1,30; 11,90]	43	5,50 [2,00; 12,90]	0.147
17	INR	35	1,10 [0,86; 1,44]	44	1,13 [0,86; 1,49]	0.730
18	pH	16	7,38 [7,31; 7,55]	26	7,40 [7,20; 7,56]	0.578
19	pCO ₂	16	32,20 [22,30; 49,00]	26	33,35 [24,50; 50,00]	0.214
20	pO ₂	16	116,00 [27,00; 192,00]	26	125,00 [70,20; 178,00]	0.707
21	SO ₂	16	98,85 [96,90; 100,00]	26	95,20 [35,00; 98,90]	0.000
22	HCO ₃ -	16	21,70 [16,40; 27,10]	26	21,80 [16,00; 26,30]	0.796

Table 11. Descriptive statistics of signs on Day 1

No.	Sign Code	Favourable		Unfavourable		Mann-Whitney Test, p-level
		n	Median [Min.; Max.]	n	Median [Min.; Max.]	
1	BT	50	36,80 [36,00; 37,80]	45	37,20 [36,00; 39,00]	0.064
2	HR	50	79,50 [68,00; 130,00]	45	101,00 [68,00; 126,00]	0.005
3	SBP	50	120,00 [90,00; 140,00]	45	120,00 [75,00; 150,00]	0.548
4	Hb	30	94,50 [64,00; 145,00]	44	89,00 [64,00; 140,00]	0.139
5	WBC	29	9,62 [4,01; 15,44]	44	10,59 [3,01; 21,20]	0.217
6	Lym	29	1,05 [0,14; 2,63]	44	0,65 [0,17; 1,70]	0.000
7	Mono	29	0,95 [0,13; 1,87]	44	0,62 [0,01; 1,45]	0.075
8	NLR	29	5,88 [1,99; 19,29]	44	16,48 [3,94; 29,10]	0.000
9	LII	29	3,89 [1,86; 9,09]	44	8,89 [3,06; 16,80]	0.000
10	ALT	28	40,00 [14,00; 103,00]	42	32,00 [6,00; 100,00]	0.275
11	B	26	16,10 [4,30; 32,80]	34	10,15 [1,50; 31,90]	0.202
12	BG	28	5,89 [4,27; 7,27]	44	7,35 [4,20; 12,05]	0.000
13	Na+	25	136,00 [132,00; 144,00]	43	143,00 [134,00; 159,00]	0.000
14	K+	25	4,20 [3,30; 5,20]	43	3,90 [2,90; 4,80]	0.001
15	Cl-	25	108,00 [99,00; 116,00]	43	110,00 [98,00; 125,00]	0.026
16	BU	28	5,05 [1,90; 8,70]	43	6,60 [2,40; 19,60]	0.005
17	INR	26	1,08 [0,92; 1,25]	39	1,19 [0,99; 1,50]	0.010
18	pH	14	7,48 [7,33; 7,54]	27	7,39 [7,26; 7,53]	0.017
19	pCO ₂	14	31,70 [23,40; 38,00]	27	36,70 [24,50; 50,80]	0.024
20	pO ₂	14	132,00 [53,00; 198,00]	27	162,00 [118,00; 206,00]	0.008
21	SO ₂	14	98,70 [96,00; 100,00]	27	98,70 [95,00; 99,80]	0.425
22	HCO ₃ -	14	23,55 [17,50; 29,00]	27	23,60 [16,30; 28,00]	0.690

Table 12. Descriptive statistics of signs on Day 7

Sign		Favourable		Unfavourable		Mann-Whitney Test, p-level
No.	Code	n	Median [Min.; Max.]	n	Median [Min.; Max.]	
1	BT	50	36,70 [36,20; 38,10]	39	37,30 [35,10; 39,90]	0.037
2	HR	50	74,50 [68,00; 143,00]	39	102,00 [68,00; 151,00]	0.000
3	SBP	50	120,00 [100,00; 136,00]	39	111,00 [77,00; 160,00]	0.077
4	Hb	16	97,50 [77,00; 115,00]	35	91,00 [66,00; 117,00]	0.051
5	WBC	16	9,41 [6,99; 21,80]	35	12,18 [3,84; 19,58]	0.707
6	Lym	16	1,08 [0,53; 2,67]	35	0,90 [0,18; 1,74]	0.088
7	Mono	16	1,32 [0,39; 3,06]	35	0,92 [0,07; 2,52]	0.025
8	NLR	16	7,67 [2,49; 14,58]	35	11,20 [3,49; 27,52]	0.006
9	LII	16	3,57 [1,77; 7,11]	35	6,48 [2,31; 19,50]	0.000
10	ALT	17	50,00 [13,00; 157,00]	35	46,00 [5,00; 196,00]	0.762
11	B	15	23,00 [6,30; 98,40]	29	11,30 [1,70; 29,90]	0.049
12	BG	17	5,39 [3,86; 7,66]	35	6,65 [4,10; 10,00]	0.019
13	Na+	17	137,00 [131,00; 143,00]	34	143,45 [129,00; 152,00]	0.000
14	K+	17	4,40 [3,60; 5,20]	34	3,90 [2,80; 5,00]	0.003
15	Cl-	17	106,00 [97,00; 115,00]	33	111,00 [94,00; 123,00]	0.002
16	BU	17	4,40 [2,10; 8,70]	34	9,10 [3,20; 19,20]	0.003
17	INR	16	1,23 [1,02; 1,45]	33	1,28 [0,96; 1,59]	0.400
18	pH	10	7,46 [7,36; 7,52]	21	7,40 [7,23; 7,52]	0.079
19	pCO ₂	10	32,00 [25,00; 43,50]	21	33,40 [28,20; 38,90]	0.540
20	pO ₂	10	82,50 [42,50; 150,00]	21	148,00 [48,00; 212,00]	0.006
21	SO ₂	10	98,50 [95,00; 99,60]	21	98,30 [95,00; 100,00]	0.767
22	HCO ₃ ⁻	10	23,45 [18,30; 28,40]	21	21,60 [14,70; 28,50]	0.123

Table 13. Descriptive statistics of signs on Day 14

Sign		Favourable		Unfavourable		Mann-Whitney Test, p-level
No.	Code	n	Median [Min.; Max.]	n	Median [Min.; Max.]	
1	BT	36	36,60 [36,20; 37,50]	21	37,40 [36,00; 39,20]	0.000
2	HR	36	77,00 [68,00; 95,00]	21	98,00 [57,00; 144,00]	0.000
3	SBP	36	120,00 [112,00; 135,00]	21	100,00 [60,00; 138,00]	0.000
4	Hb	11	96,00 [80,00; 124,00]	17	86,00 [81,00; 112,00]	0.121
5	WBC	11	12,24 [5,77; 21,04]	17	12,84 [7,51; 22,61]	0.925
6	Lym	11	1,18 [0,65; 2,07]	17	0,83 [0,28; 1,76]	0.086
7	Mono	11	0,93 [0,43; 2,62]	17	0,63 [0,38; 1,19]	0.024
8	NLR	11	7,96 [2,75; 16,29]	17	12,96 [4,84; 31,32]	0.060

Continuation of Table 13

9	LII	11	5,57 [2,17; 12,27]	17	9,68 [3,41; 13,27]	0.038
10	ALT	14	47,50 [12,00; 95,00]	16	65,30 [8,00; 177,00]	0.350
11	B	10	14,45 [5,20; 47,50]	14	8,80 [2,70; 25,20]	0.396
12	BG	14	5,97 [4,21; 7,07]	14	7,20 [4,80; 14,09]	0.018
13	Na+	9	138,00 [134,00; 139,00]	15	142,00 [132,00; 163,00]	0.056
14	K+	9	4,90 [3,80; 6,20]	15	4,10 [3,00; 5,20]	0.025
15	Cl-	9	105,00 [94,00; 114,00]	15	115,00 [94,00; 127,00]	0.152
16	BU	13	5,50 [2,00; 13,20]	16	7,25 [3,30; 21,20]	0.105
17	INR	10	1,32 [1,09; 1,44]	16	1,51 [1,18; 1,81]	0.003
18	pH	4	7,45 [7,43; 7,50]	10	7,39 [7,24; 7,44]	0.009
19	pCO ₂	4	33,10 [28,80; 37,20]	10	36,25 [22,30; 51,00]	0.525
20	pO ₂	4	75,40 [49,20; 135,00]	10	132,00 [43,00; 148,00]	0.437
21	SO ₂	4	97,60 [85,50; 99,00]	10	94,45 [80,00; 98,70]	0.358
22	HCO ₃ -	4	24,50 [22,50; 25,80]	10	23,25 [14,00; 26,60]	0.396

Table 14. Descriptive statistics of signs for Day -1

Sign		Unfavourable			
No.	Code	n	Median [Min.; Max.]	Lower quartile	Upper quartile
1	BT	41	37,30 [36,10; 40,60]	36.80	38.70
2	HR	44	101,50 [56,00; 155,00]	87.00	119.00
3	SBP	44	82,50 [50,00; 117,00]	70.00	92.50
4	Hb	36	84,50 [61,00; 117,00]	79.50	95.00
5	WBC	36	11,91 [2,43; 25,20]	8.28	16.35
6	Lym	36	0,81 [0,21; 1,98]	0.50	1.22
7	Mono	36	0,69 [0,03; 1,99]	0.33	1.34
8	NLR	36	12,00 [3,52; 39,31]	8.49	22.80
9	LII	36	7,83 [3,30; 26,24]	5.18	13.70
10	ALT	36	40,50 [9,00; 100,00]	22.50	59.00
11	B	36	10,85 [2,00; 29,10]	7.80	18.35
12	BG	35	7,79 [3,85; 17,57]	5.60	11.20
13	Na+	36	147,00 [131,00; 168,00]	141.50	153.00
14	K+	36	4,00 [2,70; 5,90]	3.35	4.60
15	Cl-	36	114,00 [96,00; 132,00]	107.00	119.00
16	BU	36	12,65 [2,60; 40,30]	6.60	20.45
17	INR	34	1,50 [1,07; 2,18]	1.31	1.74
18	pH	23	7,37 [7,10; 7,52]	7.27	7.44
19	pCO ₂	23	34,50 [17,00; 57,00]	29.20	48.70
20	pO ₂	23	87,50 [40,20; 299,00]	59.00	182.00
21	SO ₂	23	95,20 [80,80; 99,20]	88.00	98.50
22	HCO ₃ -	23	20,20 [11,10; 27,60]	18.30	23.20

CHAPTER 4.

CLINICAL AND BIOMETRIC APPROACH TO ASSESSING THE SEVERITY OF THE PHYSIOLOGICAL STATUS OF PATIENTS WITH CONCOMITANT MIDFACE TRAUMA

Decompensation of the physiological status of the patients is a complex and heterogeneous process, which is based on severe violations of the function of vital organs and body systems. The pattern of these violations may differ among different patients, depending on which systems and organs are more affected. The main element in the decompensation of the physiological status can be severe violations of the functions of one vital system or one or more organs. It is this, as well as the difference between the initial (before the injury) state and the physiological resources of the body in different patients, that causes the heterogeneity of this process.

To analyse the patterns of changes in signs reflecting the functional state of the organism, as well as the correlation relationships of these signs, we used indicators of signs collected at the fifth time point (on the last day before the unfavourable outcome).

The total number of patients in the first data subset with an unfavourable outcome was 45. The number of patients included in the study at the fifth time point (-1 day), in whom all signs (22) were measured, was 23.

In order to avoid significant data loss while reducing the number of signs and to maintain the maximum level of informativeness, factor analysis (the method of principal components (PCA)) was used, which is based on the correlation of signs. The nature of the distribution of the values of the sign 21/SO₂ is more like categorical and binary signs, despite the attempt to increase the symmetry of its values using the logarithm procedure. For this reason, the sign was excluded from this stage of biometric analysis.

To address the issue of the number of factors that need to be identified as the main ones, we were guided by the criterion of interpretability and invariance, which assumes the possibility of identifying factors as long as they can be interpreted. At the

same time, we took into account the contribution of each factor (eigenvalues) and its share of the total accumulated share of all factors.

The evaluation of the interpretability of factors and their contributions allowed us to identify three factors. The number of signs in the selected factors was reduced based on their factor loads. Signs with a factor load of <0.6 or >-0.6 are excluded from the factors. At that, the total number of signs decreased from 22 to 11. The results of the confirmatory factor analysis are presented in Table 15.

Table 15. Results of confirmatory factor analysis of signs at the fifth time point (Day -1)

Sign		Factors		
No.	Code	1	2	3
4	Hb	-0.04	-0.83	-0.11
6	Lym	-0.92	0.06	0.13
7	Mono	-0.86	0.10	0.07
8	NLR	0.84	0.01	-0.34
9	LII	0.94	-0.02	-0.13
13	Na+	0.30	0.14	0.81
15	Cl-	0.14	0.16	0.87
16	BU	0.26	0.01	0.84
18	pH	0.08	0.91	-0.14
19	pCO ₂	-0.07	-0.85	0.08
22	HCO ₃ ⁻	-0.09	0.88	-0.20
Factor contribution		3.38	3.07	2.36
The share of the factor's contribution, %		31	28	21
Accumulated share, %		31	59	80

Table 16 shows the correlations of the signs on the last day before the unfavourable outcome and the structure of the correlation pleiades for each of the three factors.

Table 16. Correlations of the signs of the three identified factors

Sign	Lym	Mono	NLR	LII	Na+	Cl-	BU	Hb	pH	pCO2	HCO3-
Lym	1.00	0.72	-0.73	-0.80	-0.11	-0.04	-0.22	0.03	-0.07	0.00	0.11
Mono	0.72	1.00	-0.52	-0.75	-0.17	-0.14	-0.14	-0.05	0.03	0.06	0.13
NLR	-0.73	-0.52	1.00	0.82	-0.07	-0.18	0.05	0.00	0.12	-0.02	-0.03
LII	-0.80	-0.75	0.82	1.00	0.09	0.01	0.10	0.06	0.06	-0.07	-0.05
Na+	-0.11	-0.17	-0.07	0.09	1.00	0.64	0.50	-0.09	0.00	0.02	0.04
Cl-	-0.04	-0.14	-0.18	0.01	0.64	1.00	0.64	-0.18	0.02	-0.18	-0.06
BU	-0.22	-0.14	0.05	0.10	0.50	0.64	1.00	-0.14	-0.02	0.04	-0.25
Hb	0.03	-0.05	0.00	0.06	-0.09	-0.18	-0.14	1.00	-0.71	0.51	-0.64
pH	-0.07	0.03	0.12	0.06	0.00	0.02	-0.02	-0.71	1.00	-0.75	0.75
pCO2	0.00	0.06	-0.02	-0.07	0.02	-0.18	0.04	0.51	-0.75	1.00	-0.71
HCO3-	0.11	0.13	-0.03	-0.05	0.04	-0.06	-0.25	-0.64	0.75	-0.71	1.00

Figure 4 shows a cross-section diagram of the correlation cylinder (correlation pleiades of P.V. Terentyev) of the signs for the last day before the onset of an unfavourable outcome. In general, the diagram illustrates the results of the factor analysis well.




Figure 4. A cross-section diagram of the correlation cylinder at the levels $|r| = 0.5$ and $|r| = 0.7$ (for patients with an unfavourable outcome on the last day before the outcome). The thickness of the lines is proportional to $|r|$ [cited by: 35, p. 15].

The selected three factors can be considered as a 'macro portrait' that characterises the physiological status of the patient on the last day before the onset of an unfavourable outcome. The interpretation of these factors is given below:

1. The first factor is represented by four signs: Leukocyte Intoxication Index (LII), Neutrophil-Lymphocyte Ratio (NLR), absolute number of lymphocytes (Lym) and monocytes (Mono). This factor reflects the nature of the morphological relationships of the main cells of the immune system and the severity of general intoxication of the body and infectious complications. The frequency and distribution of infectious complications in the studied group of patients are given in Table 17.

Table 17. The frequency of infectious complications in the studied group of patients (n=23)

Indicator	Number of patients, n=23	
	absolute number	%
Infectious complications	23	100
Pneumonia	21	91.3
Sepsis, severe sepsis, septic shock	11	47.8
Meningitis, meningoencephalitis	8	34.8
Sinusitis, polysinusitis	4	17.4
Cystopyelonephritis	2	8.7
Peri-mandibular phlegmons	2	8.7
Tracheobronchitis, endobronchitis	1	4.3
Other infectious complications	2	8.7

2. The second factor includes four signs: blood acidity level (pH), partial pressure of carbon dioxide in arterial blood (pCO₂), blood bicarbonate level (HCO₃⁻), and haemoglobin level (Hb). The second factor reflects the acid-base state and the adequacy of pulmonary ventilation and tissue oxygenation.

Depending on the type, severity, and origin of the violation of the acid-base state, it is possible to determine the nature of functional disorders of the body systems that provoke this violation. A violation of the acid-base state of metabolic origin indicates the intensity of the process of tissue destruction, while the respiratory origin of these disorders indicates a violation of pulmonary ventilation and tissue oxygenation.

Acute acidosis (pH <7.35) was observed in 9 out of 23 patients (39.1%). Mixed acute acidosis (respiratory (pCO₂ >48 mmHg) and metabolic (HCO₃⁻ <22.2 mmol/L)) was observed in 5 out of 9 patients (55.6%). Metabolic acute acidosis was observed in 4 patients (44.4%).

Acute alkalosis (pH >7.45) was observed only in 4 out of 23 patients (17.4%). A mixed type of acute alkalosis (respiratory (pCO₂ <32 mmHg) and metabolic (HCO₃⁻ >28.3 mmol/L)) was observed in one patient (25%). Respiratory type of acute alkalosis was observed in three patients (75%).

Thus, violations of the acid-base state (acidosis or alkalosis) were observed in 13 out of 23 patients (56.5%). These violations indicate severe metabolic disorders in

the body of the patients in 10 out of 13 (76.9%) cases and respiratory disorders in 9 cases (69.2%) (a mixed type of acid-base disorder was found in 6 patients).

In addition to the acid-base state component, there is also a respiratory component in the second factor. The main causes of hypoxemia are anemia and impaired blood oxygenation in the lungs due to a decrease in pO_2 . Anemia in the last day before the outcome occurred in all the patients. Mild anemia (haemoglobin = 90-119 g/l) was noted in 9 out of 23 patients (39.1%). Moderate anemia (haemoglobin = 70-89 g/l) - in 13 (56.5%); and severe anemia (haemoglobin <70 g/l) - in one patient (4.3%).

Hypoxemia ($pO_2 < 75$ mmHg) was observed in 9 out of 23 patients (39.1%), hyperoxemia ($pO_2 > 120$ mmHg) - in 10 (43.5%). It should be noted that hyperoxemia occurs as a result of intensive oxygen therapy, which is carried out in order to increase the saturation of haemoglobin with oxygen. However, in patients with anemia, this approach does not always lead to an improvement in SO_2 , and only to an increase in the concentration of oxygen dissolved in plasma. Respiratory failure of the first type ($pO_2 < 60$ mmHg and $pCO_2 < 50$ mmHg) was noted in five patients, and of the second type ($pO_2 < 60$ mmHg and $pCO_2 > 50$ mmHg) only in one. In general, respiratory system dysfunction was observed in 19 out of 23 patients (82.6%).

3. The third factor is represented by three signs: the concentration of sodium ions (Na^+), chlorine ions (Cl^-), and the level of urea (BU) in the blood. This factor reflects the functional state of the excretory system and the state of water-electrolyte metabolism.

Hypernatremia ($Na^+ > 145$ mmol/L) was observed in 17 out of 23 patients (73.9%); and hyponatremia ($Na^+ < 136$ mmol/L) - in three (13%). Hyperchloremia ($Cl^- > 107$ mmol/L) was observed in 17 patients (73.9%); and hyperchloremia - only in one (4.3%). An increase in the concentration of urea in the blood above the physiological norm (9.2 mmol/L) was observed in 13 patients (56.5%). It should be noted that hypovolemia and/or excessive infusion therapy with sodium chloride solutions could be the causes of impaired concentrations of sodium and chlorine ions in the blood. For

this reason, changes in the concentration of Na⁺ and Cl⁻ in the blood are considered indirect signs of impaired function of the excretory system. However, the direct correlation of the concentrations of these ions with the level of urea in the blood ($r(\text{Na}^+) = 0.5$; $r(\text{Cl}^-) = 0.64$) allows us to conclude that these violations are associated, among other things, with a violation of the function of the excretory system.

After the allocation of three factors, the 'scores' of each factor were calculated for the patients of the first subset of observations at each of the time points (days 1, 3, 7, and 14). The calculation of factor scores by this approach allows us to study the selected factors in dynamics. It is known that the dynamics of the average values of signs and average variances reflect the nature of the course of the pathological process [39, 46]. Based on the results of a comparative analysis of the scores of a certain factor in dynamics among patients with favourable and unfavourable outcomes, it is possible to assess the significance of this factor (and the signs forming it) for assessing the severity of the physiological status of the patients in the observation subset. The number of patients whose factor scores were calculated in dynamics is presented in Table 18. Only patients treated in the ICU wards at each of the time points were included in the analysis at this stage.

Table 18. The number of patients when calculating the scores of factors in dynamics

Time point	Number of patients					
	Total		Favourable outcome		Unfavourable outcome	
	absolute	%	absolute	%	absolute	%
Day 1	42	100	16	38.1	26	61.9
Day 3	41	100	14	34.1	27	65.9
Day 7	31	100	10	32.3	21	67.7
Day 14	14	100	4	28.6	10	41.4

Figures 5, 6, and 7 show the dynamics of the scores of the selected three factors on days 1, 3, 7, and 14 in patients with favourable and unfavourable outcomes, as well as on the last day before the outcome in patients with an unfavourable outcome.

Visual analysis of Figure 5 shows that the average values of the scores of the first factor in the group of patients with an unfavourable outcome are greater than those

in the group of patients with a favourable outcome at all time points. The variance of the scores of the first factor on the first day is significantly greater in the group of patients with an unfavourable outcome. The greatest variance of the scores of the first factor is observed in patients with an unfavourable outcome on the last day before the onset of an unfavourable outcome.




Figure 5. Diagram of the dynamics of the scores of the first factor.

In Figure 6, it can be seen that the differences in the average values of the scores of the second factor in the groups are insignificant and most pronounced on the first day after the injury. The variance of the scores of the second factor on the first and seventh day in the group of patients with an unfavourable outcome is greater than those in the group of patients with a favourable outcome. On the third day, the variance of the scores of the second factor is slightly higher in patients with a favourable outcome. The greatest variance of the scores of the second factor is observed on the last day before the outcome in the patients with an unfavourable outcome.




Figure 6. Diagram of the dynamics of the scores of the second factor.

Differences in the average values of the scores of the third factor in the groups are most pronounced on the third and seventh days. These differences increase on the third and seventh days and decrease on the fourteenth. The variance of the scores of the third factor in the group of patients with an unfavourable outcome on the first and seventh days is greater than that in the group of patients with a favourable outcome. As well as the variance of the scores of the first and second factors, the greatest variance of the scores of the third factor is observed in the group of patients with an unfavourable outcome on the last day before the onset of an unfavourable outcome (Figure 7).




Figure 7. Diagram of the dynamics of the scores of the third factor.

To test the hypothesis of equality of the mean values of the factor scores in the groups, Student's t-test for independent groups was used. The presence of statistically significant differences in the mean values of the first factor on the third ($p= 0.000$), seventh ($p= 0.013$), and fourteenth days ($p= 0.029$) was established. It was also found that there were statistically significant differences in the average values of the third factor only on the seventh day ($p= 0.009$). In all other cases, the differences were statistically insignificant. These results are presented in Table 19.

Table 19. Comparative analysis of the average values of factor scores in patients with favourable and unfavourable outcomes

Factor	Time point (p-level)			
	Day 1	Day 3	Day 7	Day 14
Factor 1	0.190	0.000	0.013	0.029
Factor 2	0.116	0.227	0.512	0.564
Factor 3	0.398	0.104	0.009	0.228

Based on a visual analysis of the dynamics diagrams of scores of the selected factors, as well as a comparative analysis of the average values of the scores of the factors, it can be concluded that the separation of the groups of the first subset of observations, depending on the nearest outcomes, is most clearly based on the first factor.

Further analysis was aimed at studying the signs forming the first factor (LII, NLR, Mono, Lym) separately. Table 20 shows data on the number of patients in whom these signs were measured at each of the time points.

Table 20. The number of patients in the comparative analysis of the signs making the first factor

Time point	Number of patients					
	Total		Favourable outcome		Unfavourable outcome	
	absolute	%	absolute	%	absolute	%
Day 1	86	100	44	51.2	42	48.8
Day 3	73	100	29	39.7	44	60.3
Day 7	51	100	16	31.4	35	68.6
Day 14	28	100	11	39.3	17	60.7
Day -1	36	100	-	-	36	100

Figure 8 shows a diagram of the dynamics of absolute numbers of lymphocytes. Statistically significant differences in the mean values of this sign among the patients with favourable and unfavourable outcomes were observed on Day 1 ($p=0.001$) and 3 ($p=0.000$). On Days 7 and 14, the differences in averages were statistically insignificant ($p=0.086$ and 0.082 , respectively). The greatest variance of the values of the Lym sign is observed on Day 1.




Figure 8. Diagram of dynamics of absolute numbers of lymphocytes.

Figure 9 shows a diagram of the dynamics of absolute numbers of monocytes. The differences in the average values of this sign by comparison groups turned out to be statistically insignificant on Days 1 ($p=0.141$) and 3 (0.074) days; and significant on days 7 and 14 ($p=0.024$ and 0.022 , respectively). The greatest variance in the absolute number of monocytes can be seen on Day 7 after the injury.




Figure 9. Diagram of the dynamics of absolute numbers of monocytes.

Figure 10 shows a diagram of the dynamics of the values of the Neutrophil-Lymphocyte Ratio. Statistically significant differences in the mean values of this sign among the patients with favourable and unfavourable outcomes were observed on Day 1 ($p=0.012$), 3 ($p=0.000$), and 7 ($p=0.005$). On Day 14, the differences in means were statistically insignificant ($p=0.059$). The greatest variance in the values of the NLR sign is observed on Day 1 in the group of patients with an unfavourable outcome.




Figure 10. Diagram of the dynamics of the values of the Neutrophil-Lymphocyte Ratio.

A diagram of the dynamics of values of the leukocyte intoxication index is shown in Figure 11. Differences in the average values of this sign by comparison groups turned out to be statistically significant at all time points during the course of the traumatic disease. The greatest variance in the values of the LII sign is observed on the last day before the unfavourable outcome. The results of testing the hypothesis of the equality of the mean values of the signs of Lym, Mono, NLR, and LII among the patients with favourable and unfavourable outcomes using the Mann-Whitney test are presented in Table 21.




Figure 11. Diagram of the dynamics of values of the leukocyte intoxication index.

Table 21. The results of testing the hypothesis about the equality of the mean values of the signs of Lym, Mono, NLR, and LII in patients with favourable and unfavourable outcomes

Time point	Sign		Mann-Whitney test, (p=)	Time point	Sign		Mann-Whitney test, (p=)
	No.	Code			No.	Code	
Day 1	6	Lym	0.001	Day 7	6	Lym	0.086
	7	Mono	0.141		7	Mono	0.024
	8	NLR	0.012		8	NLR	0.005
	9	LII	0.029		9	LII	0.000
Day 3	6	Lym	0.000	Day 14	6	Lym	0.082
	7	Mono	0.074		7	Mono	0.022
	8	NLR	0.000		8	NLR	0.059
	9	LII	0.000		9	LII	0.037

Thus, based on the comparative analysis carried out, it can be concluded that the separation of the groups of the subset of observations depending on the immediate

outcomes is most clearly determined by the first factor, which includes the following four signs: absolute numbers of lymphocytes and monocytes, NLR, and LII. This factor reflects the nature of the morphological relationships of the main cells of the immune system and the severity of general intoxication of the body and infectious complications. Further analysis of the signs forming the first factor showed that statistically significant differences in the mean values among the patients with favourable and unfavourable outcomes simultaneously on days 1, 3, 7, and 14 were observed only in the LII sign.

CHAPTER 5.

METHODOLOGY FOR PREDICTING THE IMMEDIATE OUTCOMES OF
COMCOMITANT MIDFACE TRAUMA

The currently existing methods and scales for predicting the outcome of injury have been developed to solve one or more specific tasks. Most often, these methods were intended to solve the problem of sorting the patients and determining the order of medical care at the pre-hospital or hospital stages. In such methods, the prognosis is calculated at the earliest possible time or during the first day after the injury.

In making the design of this study, the task of triage was not set, and the first question that was planned to be answered was what time point could be considered critical and therefore 'optimal' for making predictions, and at what time predicting the immediate outcome of this type of injury in general becomes possible. To solve this issue, a method of statistical analysis was applied - factor analysis (PCA) with the inclusion of a set of 22 physiological signs mentioned in the previous chapter, as well as the 'outcome' sign characterising the group affiliation of the patients (favourable or unfavourable outcomes). This analysis was carried out in a separate form with data collected on Days 1, 3, 7, and 14.

The conducted factor analysis showed that on Days 1, 7, and 14 the sign of group association 'outcome' got included in two or more factors. At that, the maximum factor load was 0.74. On Day 3, this sign was included only in the first factor with a factor load of 0.84. Based on these results, it can be concluded that, taking into account a set of physiological signs included in our work, the most optimal time point (critical point) for predicting the immediate outcomes of a concomitant midface trauma is the third day after the injury. The results of factor analysis with the values of signs for the third day are presented in Table 22. The number of patients included in this analysis on the third day was 36.

Table 22. Results of factor analysis on the third day after the injury

Sign		Factor							
No.	Code	1	2	3	4	5	6	7	8
-	Outcome	-0.84	0.31	-0.01	-0.1	-0.17	0.12	-0.18	0.14
1	BT	-0.19	-0.53	0.08	0.33	-0.25	-0.27	0.26	0.1
2	HR	-0.02	-0.04	-0.04	-0.27	-0.06	-0.84	-0.06	-0.1
3	SBP	-0.2	-0.03	-0.2	0.29	-0.01	-0.02	-0.63	-0.27
4	D	-0.34	-0.48	0.24	-0.45	0.08	0.32	0.16	-0.18
5	WBC	0.02	-0.5	0.44	0.29	0.59	0.04	0.1	0.11
6	Lym	-0.61	0.25	0.56	0.23	0.33	0.12	0.06	-0.12
7	Mono	-0.55	-0.17	0.37	0.45	0.34	-0.24	-0.02	0.01
8	NLR	0.61	-0.66	-0.29	-0.06	0.13	-0.05	-0.05	0.18
9	LII	0.71	-0.53	-0.23	-0.2	0.11	0.1	-0.03	0.14
10	ALT	-0.18	-0.22	-0.59	0.32	-0.34	0.19	0.26	-0.12
11	B	-0.44	-0.67	-0.02	0.11	-0.19	0.3	0.09	-0.09
12	BG	0.68	-0.05	0.21	0.3	-0.01	0.06	-0.06	-0.38
13	Na+	0.7	0.22	0.34	0.38	-0.03	0.15	0.17	0.07
14	K+	-0.46	-0.04	-0.23	-0.2	0.31	0.33	-0.29	0.34
15	Cl-	0.62	0.29	0.21	0.45	-0.13	0.09	-0.08	0.34
16	BU	0.29	-0.2	-0.02	-0.17	0.54	-0.19	-0.28	0.14
17	INR	0.48	0.03	-0.18	-0.04	0.35	0.16	-0.11	-0.62
18	pH	-0.58	0	-0.45	0.32	0.22	-0.17	0.28	-0.08
19	pCO2	0.6	0.44	0.15	-0.36	0.07	0.12	0.25	0.03
20	pO2	0.49	0.13	-0.48	0.38	0.12	-0.06	0.08	0.07
21	SO2	-0.16	0.32	-0.66	0.32	0.37	0.14	-0.06	0.08
22	HCO3-	-0.16	0.31	-0.29	-0.25	0.5	-0.08	0.53	-0.03
Accumulated share, %		24	36	47	56	64	70	75	80

The data used in the development of the methodology for predicting immediate outcomes are as follows: indicators of 22 physiological signs, the age of the patients, the severity of injuries on the scale of military field surgery - Injury (Mechanical Trauma), data on the development of cerebral oedema (CE) in the patients after injury and on their history of diseases of the cardiovascular system (CVD). The development of the forecasting methodology was carried out using the statistical method of discriminant analysis (DA).

Among the 22 physiological signs, signs of blood gas composition (SO₂, pH, pCO₂, HCO₃⁻, pO₂) were measured much less than others. This is due to the fact that this analysis was performed in intensive care wards, the high cost of this analysis, and various technical difficulties. For this reason, as well as to increase the availability of the developed forecasting methodology, discriminant analysis was carried out in two versions: with the inclusion of signs of blood gas composition and without them.

5.1. The First Variant of the Discriminant Function

The first variant of the discriminant analysis included the following signs: all 22 physiological signs (including signs of blood gas composition), the age of the patient, the severity of injuries on the MFS - I (MT) scale, the presence or absence of CVD in the anamnesis, and the development of CE after injury.

The step-by-step DA procedure (with inclusion) left 9 out of 26 signs. The first variant of DA included 40 patients. The classification in the sample was 100% (without inversions). Table 23 shows the factor loads of the discriminant function (DF) features. The classification of patients and the values of DF are shown in Figure 12.

Table 23. Factor loads of DF (first variant) [cited by: 36, p. 36].

Sign	Factor loads of DF
CE	-0.47
LII (on Day 3)	0.37
pO ₂ (on Day 3)	0.25
BG (on Day 3)	0.24
K+(on Day 3)	-0.24
INR (on Day 3)	0.21
pCO ₂ (on Day 3)	0.18
CVD	-0.18
Age	0.16




Figure 12. Histogram of DF values (first variant).

Calculation of the forecast of the nearest outcomes in the first variant of the DF is carried out according to the following methodology:

1. Calculation of the discriminant function by the formula:

$$DF = 0.01432*age - 2.16839*CE - 0.76152*CVD + 0.13033*LII + 0.15547*BG - 1.12332*K(+) + 1.47946*INR + 0.04215*pCO_2 + 0.01482*pO_2 + 0.66882 \quad (3)$$

where CE – if the patient developed CE - 1 in the patient; in the absence of CE - 2.

CVD – if there is a history of CVD - 1; if there is none - 2.

Indicators of the following signs were used for calculation: LII, pO₂, BG, K⁺, INR, pCO₂, measured on the third day after the injury.

2. Statistical decisive rule: if $DF < 0$, then the prognosis for the patient is considered favourable; if $DF > 0$ – unfavourable.

Among the signs included in the first variant of DF, the highest factor load was observed in the sign of CE (Table 17). With the development of such a severe

complication in the patient, the probability of a fatal outcome seriously increases. The probability of a fatal outcome also increases with an increase in the values of the signs of LII, pO₂, BG, INR, pCO₂ and with a decrease in the values of K⁺ (measured on the third day after injury). The lowest factor loads were observed in signs of age and CVD. With an increase in the age of the patients and if they have a history of CVD, the probability of a fatal outcome also increases.

5.2. The Second Variant of the Discriminant Function

The second variant of the discriminant analysis included the following signs: 17 physiological signs (not included signs of blood gas composition: SO₂, pH, pCO₂, HCO₃⁻, pO₂), the age of the patient, the severity of damage on the MFS - I (MT) scale, the history of CVD in the anamnesis, and the development of CE after injury.

Step-by-step DA procedure (with inclusion) left 8 signs: WBC (on Day 1), log LII (on Day 3), K⁺ (on Day 3), log BU (on Day 3), log Lym (on Day 3), the development of CE, age, and CVD. The number of patients included in the analysis was 64. The classification in the sample was 98.4% (1 inversion). Factor loads of DF signs are presented in Table 24. The classification of the patients and the values of the DF are shown in Figure 13.

Table 24. Factor loads of the discriminant function (second variant)
[cited by: 36, p. 37].

Sign	Factor loads of DF
Cerebral oedema	-0.49
log LII (on Day 3)	0.41
K ⁺ (on Day 3)	-0.25
CVD	-0.24
log BU (on Day 3)	0.22
Age	0.22
log Lym (on Day 3)	-0.22
WBC (on Day 1)	-0.10




Figure 13. Histogram of DF values (second variant).

Calculation of the forecast of the nearest outcomes in the first variant of the DF is carried out according to the following methodology:

1. Calculation of the discriminant function by the formula:

$$DF = 4.86639 \cdot \log LII (\text{on Day 3}) - 1.01252 \cdot K^+ (\text{on Day 3}) + 2.64833 \cdot \log Lym (\text{on Day 3}) + 2.43754 \cdot \log BU (\text{on the Day 3}) - 0.05692 \cdot WBC (\text{on Day 1}) - 0.01599 \cdot \text{age} - 2.32129 \cdot CE - 1.18639 \cdot CVD + 6.68479 \quad (4)$$

where CE – if the patient developed CE - 1; in the absence of CE - 2.

CVD – if there is a history of CVD - 1; if there is none - 2.

Indicators of signs were used for calculation: WBC - measured on Day 1 after injury; signs of Lym, LII, K⁺, BU - on Day 3 after injury.

2. Statistical decisive rule: if $DF < 0$, then the prognosis for the patient is considered favourable; if $DF > 0$ – unfavourable.

As in the first variant of the DF, the highest factor load among the signs included in the second variant of the DF was found in the sign of CE (Table 18). These data

indicate a serious impact that the development of such a complication has on the immediate outcomes of a concomitant midface trauma. The probability of a fatal outcome increases with an increase in the values of signs of age, LII (Day 3) and BU (Day 3), as well as with a decrease in the values of signs of WBC (Day 1), Lym, and K + (Day 3, respectively). If the patients have a history of CVD, the probability of a fatal outcome increases.

The presented forecasting technique can be used as an auxiliary tool in determining the timing of reconstructive operations. According to the theory of traumatic illness, the primary stabilisation of the physiological status of the patients occurs on the second and third days after the injury. In case of an unfavourable prognosis, it is recommended to refrain from performing reconstructive operations until the onset of stable stabilisation of the physiological status of the patient. With a favourable prognosis and a compensated physiological status of the patient, the decision to perform reconstructive operations at an early date can be considered justified.

CHAPTER 6.

TREATMENT OF PATIENTS WITH CONCOMITANT MIDFACE TRAUMA

6.1. Tactics of Multi-Stage Surgical Treatment of Patients with Concomitant Midface Trauma

The treatment of patients with severe and extremely severe combined injuries of the first and second groups of observed patients was carried out using the MST tactics, taking into account the severity and level of compensation of the physiological status of the patients.

The MST tactic in patients with concomitant midface trauma includes the following stages:

Stage I of MST corresponds to the first period of TD - the period of acute impairment of vital functions. This period lasts up to 12 hours from the moment the patient is admitted to hospital. The goal of the first stage is to eliminate the life-threatening consequences of trauma, including the elimination of asphyxia and performing tracheostomy, while creating physiological status for intensive therapy, stopping bleeding, eliminating compression of the brain; temporary fixation of fragments by orthopaedic method in order to stop bleeding, and performing elements of primary soft tissue plasty. These measures are carried out in the anti-shock operating room synchronously with similar actions of specialists of other profiles [11, 24, 25, 56, 58, 88, 89].

Primary soft tissue plasty is performed when there are extensive penetrating wounds with soft tissue defects; when there are wounds with soft tissue defects in the eyelids, nose, auricles, and lips; when parts and organs of the maxillofacial region, such as the nose, lips, and auricles, are torn off [88].

Stage II of MST corresponds to Periods 2 and 3 of TD and lasts from 2 to 10 days. At the second stage, intensive therapy is carried out to achieve temporary stabilisation of the physiological status of the patient; measures are aimed at preventing

and treating infectious complications. The second stage can be divided into 2 parts: 1-3 days - long-term ventilator therapy, infusion-transfusion therapy (transfusion of erythroconcentrate, freshly frozen plasma, plasma-substituting solutions) and de-escalation antibiotics therapy (ABT); 4-10 days - targeted ABT taking into account the culturing results, and minimally invasive operations (rehabilitation of potential sources of infectious complications). In certain cases, staged fixation of fragments is carried out using minimally invasive extra-focal osteosynthesis [70, 71]. At this stage, an exhaustive diagnosis of damaged structures using CT is also carried out. A component of complex treatment is restoration of damaged structures to functional capacity, including the insertion of a feeding tube through a stoma or the placement of a gastrostomy [11, 24, 25, 51, 56, 58, 89].

Stage III of MST is reconstructive and restorative and corresponds to the 4th TD period. The third stage can be started after the final stabilisation of the physiological status of the patient. Surgical interventions are performed in full and extra-focal osteosynthesis is replaced by intraosseous osteosynthesis in order to achieve complete stabilisation of the fracture, eliminate aesthetic defects, and restore the bite [11, 24, 56, 58, 89].

To assess the severity and level of compensation of the physiological status of the patients, a technique used at the clinical facilities of the Department of Maxillofacial Surgery and Surgical Dentistry of St Petersburg University was used, which consists in assessing the following seven signs: data on inotropic support for the patient, systolic blood pressure, heart rate, haematocrit level, haemoglobin level, concentration of blood erythrocytes, the severity of the physiological status on the Military Field Surgery Scale - Selective Physiological status Assessment (MFS-SC). The MFS - SC scale was developed by the Doctor of Medical Sciences, Professor E.K. Gumanenko at the Kirov Military Medical Academy in 1992 [20, 26, 75]. The quantitative limits of these signs for each level of compensation of the physiological status are presented in Table 25.

Table 25. Methodology for assessing the level of compensation for the physiological status of patients with concomitant craniofacial trauma

Sign	Level of compensation of the physiological status of the patients		
	Decompensation	Subcompensation	Compensation
Inotropic support	Given	Not given	Not given
Systolic Blood Pressure (mmHg)	≤ 80	81-100	>100
HR	<60 or >140	91-140	60-90
Haematocrit (%)	20-25	26-31	>31
Haemoglobin (g/l)	60-84	85-99	>99
Red blood cells ($^{10-12}/l$)	<2.5	2,6-2,9	>2.9
MFS - SC, score	70-98	50-69	<50

When using this method, the general decompensation conditions are checked first. If there is at least one decompensation condition, the general condition is assessed as decompensated. In the absence of decompensation conditions, the subcompensation conditions are checked. In the absence of decompensation and subcompensation conditions, the overall condition is assessed as compensated.

In a decompensated patient's general condition, the extent of necessary and tolerable surgical care corresponds to Stage I of MST; in a subcompensated condition - to Stage II; and in a compensated condition - to Stage III. Thus, reconstructive operations are not always postponed for the fourth period of TD (after 10 days), and their timing is determined directly taking into account the severity of the general condition of the patient.

6.2. Treatment of Patients with Severe Concomitant Midface Trauma

6.2.1. Clinical Characteristics of Patients with Severe Concomitant Midface Trauma

When creating the study groups to solve the third task of the study, the author was guided by the classification of the severity of damage on the MFS - I (MT), according to which injuries estimated at 1.0 to 12 points are attributed to severe injuries; and at ≥ 12 points - to extremely severe. 48 patients were included in a group of patients with severe concomitant midface trauma. They were divided into two groups: the first group - included 27 patients — it is the group of the author's own observations. This group will be conventionally called the first group of observed patients; the second group included data on 21 patients collected by the retrospective method. The second group will be conventionally called the first retrospective group.

In addition to injuries to the midface zone, all the patients, both in the group of observed patients and the retrospective group, were diagnosed with injuries to other anatomical areas. Injuries in the head and, including the cerebral part of the skull, were diagnosed in 100% of the patients. Open TBI was diagnosed in 25 (92.6%) patients of the first group of observed patients and in 11 (52.4%) patients of the first retrospective group; closed TBI - in 2 (7.4%) and 10 (47.6%) patients, respectively. The second most frequent injury is to the chest area; the third most frequent is to the extremities. The distribution of injuries by anatomical areas in the patients of the first group of observed patients and the first retrospective group is presented in Table 26.

Table 26. Distribution of injuries by anatomical areas in patients with severe concomitant midface trauma

Anatomical area	Group and number of patients			
	1st retrospective group n=21		1st group of observed patients n=27	
	absolute number	%	absolute number	%
Head	21	100	27	100
Neck	1	4.8	0	0
Chest	9	42.9	5	18.5
Stomach	3	14.3	0	0
Pelvis	2	9.5	0	0
Spine	3	14.3	2	7.4
Limbs	7	33.3	6	22.2

In all the patients of the first group of observed patients, as well as in 12 (57.1%) patients of the first retrospective group, most severe injuries were localised in the head area; in 4 (19%) patients of the first retrospective group - in the chest area; in 3 (14.3%) - in the extremities; in 1 (4.8%) - in the spine and abdomen. In addition to midface injuries, the patients of the first group of observed patients were diagnosed with injuries, on average, in 1.3 anatomical areas; of the first retrospective group - in 1.9.

On admission to the clinic, all the patients (n=48) were treated in the shock ward of the emergency department and, later, in the resuscitation and intensive care units. The differences in the severity of injuries in the groups on the ISS scale and the MFS - I (MT) scale are statistically insignificant (Table 27).

Table 27. Severity of injuries in the first group of observed patients and the first retrospective group

Severity of injury, score; median [min.; max.]	Group and number of patients		Mann-Whitney p-level
	1st retrospective group n=21	1st group of observed patients n=27	
MFS - I (MT)	2,75[1,2; 10,55]	2,65 [2,1; 7,65]	0.547
ISS	21 [8; 29]	21 [9; 29]	0.716

The frequency of damage to the bone structures of the midface in the patients of the first group of observed patients is higher than in the patients of the first retrospective group, and midface soft tissue injuries were diagnosed more often in the first retrospective group. The first place among the fractures of the midface bones in both groups is occupied by fractures of the eye socket walls, and the second - by fractures of the upper jaw. On average, the patients of the first group of observed patients were diagnosed with fractures of 2.2 midface bones; the patients of the first retrospective group - of 1.8. The distribution of midface injuries by groups is presented in Table 28.

Table 28. Distribution of midface injuries in patients of both groups

Anatomical location	Group and number of patients			
	1st retrospective group n=21		1st group of observed patients n=27	
	absolute number	%	absolute number	%
Upper jaw	8	38.1	15	55.6
Eye socket	12	57.1	19	70.4
Zygomatic bone	7	33.3	11	40.7
Nose bones	6	28.6	14	51.9
Soft tissues	16	76.2	10	37

In the majority of observed patients of the first group (n=26 (96.3%)) the first variant of the course of traumatic disease was observed; the second variant was observed only in one patient (3.7%). In the first retrospective group, the first variant of the course of traumatic disease was detected in 66.7% (n=14) of patients; the second variant - in 23.8% (n=5). The third variant of the course of traumatic disease was observed only in two patients (9.5%) of the first retrospective group, and in the first group of own observations it was not detected. The distribution of patients of both groups, depending on the course of the traumatic disease, as well as the immediate outcomes of the combined injury in each variant are presented in Table 29.

Table 29. Distribution of patients according to the course of the traumatic disease and the immediate outcome of the combined injury

Indicator		Group and number of patients					
		1st retrospective group n=21			1st group of observed patients n=27		
		I	II	III	I	II	III
Absolute number of patients (%)		14; (66,7%)	5; (23,8%)	2; (9,5%)	26; (96,3%)	1; (3,7%)	0; (0%)
Severity of injury (score) - median [min; max.]	MFS - I (MT)	2,75 [1,2; 10,55]			2,6 [2,1; 8,55]		-
	ISS	21 [8; 29]			21 [9; 29]		-
Survived, absolute number (%)		14; (100%)	3; (60%)	0; (0%)	26; (100%)	1; (100%)	-
Died, absolute number (%)		0; (0%)	2; (40%)	2; (100%)	0; (0%)	0; (0%)	-

6.2.2. Results of Treatment of Patients with Severe Concomitant Midface Trauma

According to the concept of multi-stage surgical treatment (Damage Control Orthopaedics), decisions on the timing and scope of necessary and permissible surgical care for the patient are made taking into account the overall physiological status severity, as well as the nature (variant) of the course of the traumatic disease. Only if this physiological status is met, the decisions made can be considered justified.

We assessed the level of compensation of the physiological status of the observed patients of the first group (n=27) on Days 1, 3, 7, and 14 of inpatient treatment. The results are presented in Table 30. As can be seen from the table, despite the presence of severe injuries, the physiological status of almost half of the patients (55.6%) was assessed as compensated. The proportion of patients with compensated physiological status increased on Day 3, and by Day 7 was 100%, which indicates complete stabilisation of their physiological status.

Table 30. The level of compensation of the physiological status of observed patients in the first group in dynamics

Day	Number of patients		Level of compensation of the physiological status					
			Compensation		Subcompensation		Decompensation	
	absolute number	%	absolute number	%	absolute number	%	absolute number	%
1	27	100	15	55.6	10	37	2	7.4
3	27	100	22	81.5	5	18.5	0	0
7	26	96.3	26	100	0	0	0	0
14	10	37	10	100	0	0	0	0

Below is a clinical example of the implementation of a multistage surgical treatment tactic in a survivor with severe concomitant midface trauma:

Clinical example No. 1:

Patient B., 26 years old, was brought to the clinic on 08 March 2019 by an ambulance team. The injury occurred as a result of an attack with the use of physical force. After examination, the patient was diagnosed with an open craniocerebral injury. Moderate brain contusion. Fracture of the base of the skull in the anterior cranial fossa and the middle cranial fossa on both sides. Fracture of the upper jaw of the upper, middle, and lower types on both sides. Fracture of the ethmoid bone. Fracture of the supraorbital rim on the left. Bilateral Type 3 Nasoorbitoethmoidal Complex Fracture with Left Medial Canthal Ligament Avulsion. Fracture of the walls of both eye sockets. Fracture of the bones of the nose. Polyposis of the Sinuses. Bilateral zygomatic arch fracture. Lacerations on the left shoulder and right thigh. Contusions of the frontal and occipital regions. Stab-cut wound of the right chest wall.

The severity of injuries on the MFS - I (MT) scale was estimated at 2.65 points (severe), on the ISS scale - at 21 points. Upon admission to the clinic, the level of consciousness on the Glasgow Coma Scale was estimated at 12 points (severe concussion). In Figure 14, sections from the CT scan of the skull of the patient are presented, taken on the third day after the injury.

The patient, upon admission to the clinic, received immediate assistance in the shock room, synchronously with the actions of specialists from other disciplines.

Emergency aid was provided, including the cessation of nasal bleeding, relief of asphyxia, and mandibular splinting performed by maxillofacial surgeons (Dr. K.G.M. and Dr. S.E.V.)




Figure 14. CT scan of the skull of the injured patient B., 26 years old, on the 3rd day after the injury: a) anterior view; b) left lateral view; c) right lateral view; d) coronal projection.

The patient was transferred to the intensive care unit (ICU). The time of the patient's stay in the ICU was 81 hours.

In the intensive care unit, the patient received the following treatments: mechanical ventilation therapy, transfusion of fresh frozen plasma, infusion therapy, and antibiotic therapy. 'On the third day, a computed tomography scan of the damaged structures was performed, along with sanitation of the maxillary sinuses.' Considering the severity of the injuries and the favourable prognosis, tracheostomy and gastrostomy were not performed.

On the sixth day of inpatient treatment and following the final stabilisation of the patient's overall physiological status, the third stage of surgical treatment was performed - reconstructive-restorative. The extent of the surgical interventions performed is as follows: reconstruction of the cranial vault defect. Reduction and osteosynthesis of the left zygomatic bone. Reduction and osteosynthesis of the upper jaw using the lower approach. Reconstruction of the lower wall of the right orbit. The duration of the operation was 5 hours (surgeons: Dr. S.E.L., Dr. K.G.M., Dr. A.K.A.). Figure 15 shows the sections from the CT scan of the skull of the patient after the reconstructive-restorative surgeries were performed.




Figure 15. CT scan of the skull of the injured patient B., 26 years old, after the reconstructive-restorative surgeries: a) anterior view; b) left lateral view; c) coronal projection; d) axial projection.

During the postoperative period, the patient experienced an infectious complication (poly-sinusitis). The patient was discharged from the clinic in a satisfactory physiological status. The duration of hospital stay was 41 days.

The main method of fixation for fractures of the craniofacial complex in the first group of observed patients was a combined approach (intraosseous metal osteosynthesis and orthopaedic fixation methods) It was applied in 14 patients (51.9%).

In isolated form, intraosseous metal osteosynthesis was performed in 7 patients (25.9%), while the orthopaedic method was used in 3 patients (11.1%). Due to the absence of functional and aesthetic impairments, fixation of the craniofacial fractures was not performed in 3 patients (11.1%).

Complicated courses of traumatic disease were observed almost three times more frequently in the retrospective group of patients compared to the observed group. Infectious complications prevailed over non-infectious ones in the patients of both groups. The frequency of the most commonly encountered infectious and non-infectious complications in patients with severe concomitant midface trauma is presented in Table 31.

Table 31. The frequency of complications in patients with severe concomitant midface trauma

Indicator	Group and number of patients			
	1st retrospective group, n=21		1st group of observed patients n=27	
	absolute number	%	absolute number	%
Complicated course of TD	10	47.6	4	14.8
Infectious complications	8	38.1	3	11.1
Non-infectious complications	2	9.5	1	3.7
Cerebral oedema	2	9.5	0	0
Pneumonia	6	28.6	0	0
Tracheobronchitis, endobronchitis	1	4.8	0	0
Meningitis, meningoencephalitis	0	0	0	0
Organ and multiple organ failure	3	14.3	0	0
Sepsis, severe sepsis, septic shock	5	23.8	0	0
Sinusitis, polysinusitis	1	4.8	3	11.1
Cystopyelonephritis	0	0	0	0
Thrombosis, thromboembolism	1	4.8	0	0
Perimaxillary abscesses and phlegmons	0	0	3	11.1
Other complications	6	28.6	1	3.7

Severe and life-threatening complications, such as cerebral oedema, organ or multiple organ failure, pulmonary embolism, and purulent-septic complications were

observed in 7 (33.3%) patients from the first retrospective group. In the observed group of patients, no life-threatening complications were identified. These data suggest a more severe course of traumatic disease in patients of the first retrospective group.

The average duration of treatment of the observed patients of the first group in the ICU wards was 3.1 ± 1 days; of the patients of the first retrospective group - 4.9 ± 3.5 days. The average duration of hospitalisation in the observed patients of the first group was 14.2 ± 7.9 days; in the patients of the first retrospective group - 18.3 ± 8 days.

The immediate outcome of severe concomitant midface trauma turned out to be favourable in all the observed patients of the first group; in the first retrospective group - in 17 (81%) of the patients. An unfavourable outcome of the injury was observed in 4 (19%) patients of the first retrospective group.

6.3. Treatment of Patients with Extremely Severe Concomitant Midface Trauma

6.3.1. Clinical Characteristics of Patients with Extremely Severe Concomitant Midface Trauma

The cohort of patients with extremely severe concomitant midface trauma includes 57 patients. Two groups were formed from this cohort: the first group includes 21 patients — this is the group personally observed by the author. This group will be conventionally called the second group of observed patients; the second group included data on 36 patients collected by the retrospective method. The second group will be conventionally called the second retrospective group.

In addition to midface injuries, all the patients, both the second observed group and the second retrospective group, were diagnosed with injuries to one or several other anatomical areas. Injuries in the head region are diagnosed more frequently than in other anatomical areas. TBI with severe brain contusion was detected in all patients (n=57). In the observed group of patients, injuries in the limb region are the second most frequent, followed by injuries in the chest region in third place. In the retrospective group of patients, injuries in the chest region were identified more

frequently than injuries in the limb region. The distribution of injuries by anatomical areas in the patients of the second observed group and the second retrospective group is presented in Table 32.

Table 32. Distribution of injuries by anatomical areas in patients with extremely severe concomitant midface trauma

Anatomical area	Group and number of patients			
	2nd retrospective group, n=36		2nd observed group, n=21	
	absolute number	%	absolute number	%
Head	36	100	21	100
Neck	0	0	1	4.8
Chest	17	47.2	4	19
Stomach	4	11.1	1	4.8
Pelvis	2	5.6	1	4.8
Spine	3	8.3	1	4.8
Limbs	13	36.1	6	28.6

In all the patients of the second observed group of their own observations and the second retrospective group (n=57), injuries leading in severity were localised in the head area. Open TBI was diagnosed in 14 (66.7%) patients of the second observed group and in 21 (58.3%) patients of the second retrospective group; closed TBI - in 7 (33.3%) and 15 (41.7%) patients, respectively. The severity of TBI on the MFS - I (MT) scale in all cases was assessed as extremely severe (≥ 12 points). In addition to midface injuries, the patients of the second observed group of were diagnosed with injuries, on average, in 1.3 anatomical areas; of the first retrospective group - in 1.9. On admission to the clinic, all the patients (n=57) were treated in the shock ward of the emergency department and, later, in the resuscitation and intensive care units.

Table 33 shows the severity of injuries in the patients of the second observed group and the second retrospective group according to the MFS-I (MT) and the ISS scales. The differences in the severity of injuries by groups are not statistically significant ($p > 0.05$).

Table 33. Severity of injuries in the second observed and the second retrospective groups

Severity of injury, score, median [min; max.]	Group and number of patients		Mann-Whitney p-levels
	2nd retrospective group, n=36	2nd observed group, n=21	
MFS - I (MT)	13,08[12,10; 24,10]	13,70 [12,30; 20,60]	0.170
ISS	30 [26; 50]	30 [29; 45]	0.673

On average, 1.8 fractures of the midface bones were diagnosed in the patients of the second observed group; 1.3 fractures - in the patients of the second retrospective group. The number of injuries to bone structures, as well as soft tissues of the midface area in the patients of the second observed group is higher than in the patients of the second retrospective group. Among the fractures of the midface bones, fractures of the zygomatic bone take the first place in the second observed group, and fractures of the walls of the orbit take the second. In the second retrospective group, fractures of the orbit walls occupy the first place, and fractures of the upper jaw - the second. The distribution of midface injuries by groups is presented in Table 34.

Table 34. Distribution of midface injuries in patients of both groups

Anatomical location	Group and number of patients			
	2nd retrospective group, n=36		2nd observed group, n=21	
	absolute number	%	absolute number	%
Upper jaw	11	30.6	8	38.1
Eye socket	18	50	11	52.4
Zygomatic bone	9	25	13	61.9
Nose bones	9	25	6	28.6
Soft tissues	21	58.3	14	66.7

In the second observed group of patients, the third course variant of traumatic disease was observed in 47.6% (n=10) of the patients. The second course variant was present in 4 patients (19%), and the first course variant was observed in 7 patients (33.3%). The third course variant of traumatic disease was detected in the patients of the second retrospective group more often than in the patients of the second observed

group, (52.8% of the patients (n= 19)). The second course variant of traumatic disease was observed in 10 (27.8%) patients of the second retrospective group, and the first variant - in 7 (19.4%). The distribution of patients of both groups, depending on the course of the traumatic disease, as well as the immediate outcomes of the combined injury in each variant are presented in Table 35.

Table 35. Distribution of patients according to the course of the traumatic disease and the immediate outcome of the combined injury

Indicator		Group and number of patients					
		2nd retrospective group, n=36			2nd observed group, n=21		
		I	II	III	I	II	III
Number of patients absolute number; (%)		7; (19,4%)	10; (27,8%)	19; (52,8 %)	7; (33,3 %)	4; (19 %)	10; (47,6%)
Severity of injury (score) - median [min; max.]	MFS - I (MT)	13,08 [12,10; 24,10]			13,70 [12,30; 20,60]		
	ISS	30 [26; 50]			30 [29; 45]		
Survived, absolute number (%)		7; (100%)	2; (20%)	0; (0%)	7; (100%)	1; (25%)	1; (10%)
Died, absolute number (%)		0; (0%)	8; (80%)	19; (100%)	0; (0%)	3; (75%)	9; (90%)

6.3.2. Results of Treatment of Patients with Extremely Severe Concomitant Midface Trauma

Dynamic monitoring of the severity of the physiological status of the patients with extremely severe concomitant trauma of the second observed group (n=21) showed that on the first day of inpatient treatment, decompensated physiological status was noted in almost half of the patients (52.4%). On the third day, the number of such patients decreased to 42.9%, and on the seventh day increased to 45%. The proportion of patients with compensated physiological status increased from 0% on day 1 to 30.8% on day 14. These data describe one of the main features of managing patients with

extremely severe trauma - a prolonged lack of stabilisation of the physiological status. This feature emphasises special relevance of the use of the MST tactic, taking into account the dynamics of the course of traumatic disease and changes in the severity of the physiological status of patients of this group. In table 36, one can see the change in the severity of the physiological status of the patients of the second observed group in dynamics.

Table 36. Level of compensation of the physiological status of observed patients in the first group in dynamics

Day	Number of patients		Level of compensation of the physiological status					
			Compensation		Subcompensation		Decompensation	
	absolute number	%	absolute number	%	absolute number	%	absolute number	%
1	21	100	0	0	10	47.6	11	52.4
3	21	100	3	14.3	9	42.9	9	42.9
7	20	95.2	5	25	6	30	9	45
14	13	61.9	4	30.8	6	46.2	3	23.1

Below is a clinical example of the implementation of the multistage surgical treatment tactic in a survivor with extremely severe concomitant midface trauma:

Clinical example No. 2:

Patient C., 19 years old, was taken to the clinic by an ambulance team on 13 June 2019 with a gunshot wound from a traumatic weapon. After examination, the patient was diagnosed with a gunshot transcranial and facial penetrating injury involving the oral cavity, left maxillary sinus, left orbital cavity, and frontal sinus. Open traumatic brain injury. Severe brain contusion with the formation of a contusion focus in the left frontal lobe. Comminuted fracture of the mandible in the left body region. Comminuted fracture of the alveolar process of the maxilla on the left side, involving all walls of the left maxillary sinus. Comminuted fracture of the lower, lateral, medial, and upper walls of the left orbit with damage to the muscular apparatus of the left eye. Comminuted fracture of the left zygomatic bone and the walls of the frontal sinus. Multiple gunshot wounds of the oral mucosa, tongue, back and side walls of the pharynx. Foreign bodies

(fragments of bones, teeth) in the soft tissues of the oral cavity, back and side walls of the pharynx. Penetrating wound of the left eyeball. Hemophthalmos on the left. Amaurosis on the left. Bilateral pneumothorax.




Figure 16. CT scan of the skull of the injured individual, C., 19 years old, at 3 days after the trauma: a) anterior view; b) left lateral view; c) coronal projection (1); d) coronal projection (2).

The severity of injuries on the MFS - I scale (GW - for gunshot wounds) was estimated at 13.7 points (extremely severe), and on the ISS scale at 30 points. Upon admission to the clinic, the level of consciousness on the Glasgow Coma Scale was

estimated at 5 points (coma). In Figure 16, sections of the cranial CT scan of the injured individual are presented at three days after the trauma.

Upon admission to the clinic, the patient's physiological status was assessed as decompensated. The first stage of MST was implemented, including primary surgical wound management of the maxillofacial area and posterior and lateral pharyngeal walls, sanitation of the left maxillary sinus with foreign body removal. Immobilisation of the upper jaw was achieved using orthopedic methods, and stepwise fixation of the mandibular fracture was performed using an extraoral approach. Tracheostomy was also performed. The procedures were carried out by maxillofacial surgeons (S.E.V., K.G.M., A.K.A., and S.E.L.)

The patient was in the intensive care unit (ICU) for 18 days. The patient underwent mechanical ventilation therapy, fresh frozen plasma transfusion, infusion and antibiotic therapy, as well as a computed tomography of the injured anatomical structures. On 16 June 2019, pleural cavity drainage was performed. On 17 June 2019, a gastrostomy was performed. On 27 June 2019, the patient's overall physiological status was assessed as compensated, and the reconstructive-restorative stage of surgical treatment was carried out, including repositioning and osteosynthesis of the mandible and reconstruction of the orbital walls (performed by maxillofacial surgeons - S.E.L., K.G.M.). Figure 17 shows the sections from the CT scan of the skull of the patient after the reconstructive-restorative surgeries were performed.

The course of the traumatic disease was complicated by the development of endobronchitis and tracheoendobronchitis. The duration of inpatient treatment was 43 days. The patient was discharged from the clinic in a satisfactory physiological status.




Figure 17. CT scan of the skull of the injured individual C., 19 years old, after the reconstructive-restorative stage of surgical treatment: a) straight view; b) left lateral view; c) axial projection; d) coronal projection.

The fixation of fractures in the midface bones was carried out in only 9 (42.9%) of patients of the second observed group. Orthopaedic methods of fixation of fractures of the midface bones were used in 6 (28.6%) patients; the combined method - in 2 (9.5%) patients; bone metallosteosynthesis - in 1 (4.8%) patient.

The complicated course of traumatic disease in the patients of the second retrospective group was observed in 83.3% of cases; in the patients of the second

observed group - in 71.4% of cases. Most of the patients of both groups with a complicated course of traumatic disease were diagnosed with both infectious and non-infectious complications. The frequency of the most commonly encountered infectious and non-infectious complications in patients with severe concomitant midface trauma is presented in Table 37.

Table 37. The frequency of complications in patients with very severe concomitant midface trauma

Indicator	Group and number of patients			
	2nd retrospective group, n=36		2nd observed group, n=21	
	absolute number	%	absolute number	%
Complicated course of TD	30	83.3	15	71.4
Infectious complications	29	80.6	15	71.4
Non-infectious complications	28	77.8	11	52.4
Cerebral oedema	21	58.3	7	33.3
Pneumonia	28	77.8	14	66.7
Tracheobronchitis, endobronchitis	0	0	5	23.8
Meningitis, meningoencephalitis	10	27.8	5	23.8
Organ and multiple organ failure	15	41.7	3	14.3
Sepsis, severe sepsis, septic shock	16	44.4	9	42.9
Sinusitis, poisinusitis	2	5.6	2	9.5
Cystopyelonephritis	6	16.7	3	14.3
Thrombosis, thromboembolism	1	2.8	1	4.8
Perimaxillary abscesses and phlegmons	2	5.6	1	4.8
Other complications	10	27.8	5	23.8

Life-threatening complications such as cerebral oedema, organ or multiple organ failure, pulmonary embolism, and purulent-septic complications were diagnosed in 29 (80.6%) patients from the second retrospective group and in 13 (61.9%) patients of the second observed group.

The frequency of patients with a fatal outcome of extremely severe concomitant midface trauma in the second observed group was 57.1% (n=12). This indicator was

higher in the patients of the second retrospective group and amounted to 75% (n=27). The comparative characteristics of patients with an unfavourable outcome are presented in Table 38.

Table 38. Comparative characteristics of patients with an unfavourable outcome of extremely severe concomitant midface trauma

Indicator		Subset and number of patients			
		2nd retrospective group, n=36		2nd observed group, n=21	
		absolute number	%	absolute number	%
Number of fatalities		27	75	12	57.1
Time of the lethal outcome, in days	1-3	0	0	0	0
	4-6	3	11.1	1	8.3
	7-9	3	11.1	3	25
	10-12	3	11.1	4	33.3
	13-15	8	29.6	1	8.3
	16-18	1	3.7	0	0
	19-21	2	7.4	0	0
	>21	7	25.9	3	25
Average time of lethal outcome, day M±m		19,9±3,7		18,3±5,4	
Severity of injury, MFS-I (MT), score, median [min.; max.]		12,6 [12,1; 22,1]		14,9 [12,6; 20,6]	
Severity of injury, ISS, score, median [min; max.]		30 [26; 45]		29 [29; 45]	
The level of consciousness on admission by GCS, median [min.; max.]		12 [4; 15]		12 [6; 15]	

The average duration of treatment of the patients of the second observed group in the ICU wards was 12.4 ± 3 days; in the patients of the second retrospective group - 12.3 ± 1.2 days. The average duration of inpatient treatment in the patients of the second observed group was 28.3 ± 5.5 days; in the patients of the second retrospective group - 22.6 ± 3 .

CONCLUSION

The problem of severe concomitant craniofacial trauma is an important task of modern medicine due to its wide prevalence, high probability of infectious complications, and mortality. Damage to the midface area due to its anatomical proximity to the cerebral part of the skull is often combined with TBI. According to M.O. Danilevich, with multiple fractures of the midface bones, patients are diagnosed with moderate and severe TBI in 90.1% of cases [28].

111 patients with severe and extremely severe concomitant midface trauma who were treated at the clinic of the State Medical Institution Alexandrovskaya Hospital of St Petersburg in the period from 2016 to 2020 have been studied. Clinical and laboratory dynamic monitoring of the severity of the physiological status of the patients was carried out. The results of treatment of the patients have been analysed.

The first task of the study was to search for syndrome complexes to assess the severity of the physiological status of patients with concomitant midface trauma in the dynamics of the course of traumatic disease. To address this task, 22 physiological signs were measured, reflecting the degree of dysfunction of vital organs and body systems and fully characterising the physiological status of the patient. Physiological signs were examined at five time points: the 1st, 3rd, 7th, and 14th days from the moment of the injury, as well as on the last day of life of the patients with an unfavourable outcome.

This paper presents a new algorithm for biometric analysis of medical data for patients with severe trauma. The description and justification of this variant are presented below:

The measurement of the studied physiological signs was carried out in two time variants: the first variant is the time of the course of a traumatic disease (traditionally used in medical research). The time count in the first variant begins from the moment of injury, and the time points correspond to the periods of traumatic disease. The second variant of time is the reverse, and since the closest outcome of the trauma is the indicator of group affiliation in the first research task, we refer to the second variant of

time as the time of waiting for the outcome. The values of the studied signs were also collected for the last day of life of patients with a fatal outcome.

Factor analysis (PCA) was performed once with data collected on the last day of life of patients with an unfavourable outcome, and not at each time point. Next, the scores of the already identified factors were measured during the course of the traumatic disease (1st, 3rd, 7th, and 14th days).

The bottom line is that the measurement time of a feature is a key component of any medical study. The individuals who had an unfavorable outcome succumbed at different time intervals from the onset of the trauma. For this reason, time points in the course of the traumatic disease have different clinical significance for different patients. For a patient who passed away on the 4th day, the 3rd day was characterised by significant decompensation of the overall physiological status. For another patient who passed away on the 15th day, the 3rd day has a completely different clinical significance, where the physiological status may be subcompensated. For this reason, when conducting factor analysis at multiple time points during the course of traumatic disease, researchers may encounter the problem of changes in factor structure (variation in the features that form the factors). This, in turn, hinders the interpretation of the factors and prevents the study of dynamic changes in the factors over time, including the ability to conduct comparative analysis of factors among comparison groups. It is also necessary to emphasise the important clinical significance of the last days of life of patients with a fatal outcome, which are characterised by pronounced decompensation and multiple organ failure, for the problem of assessing the severity of the physiological status, which in turn supports our opinion on the feasibility of factor analysis at this time point according to the algorithm described above.

As a result of the factor analysis, three factors were identified, and the total number of signs was halved (from 22 to 11) without significant loss of informativeness. The first factor reflects the nature of morphological ratios of the main cells of the immune system and the severity of general intoxication of the body and infectious complications and includes the following signs: leukocyte intoxication index, neutrophil-lymphocyte ratio, and absolute number of lymphocytes and monocytes. The

second factor reflects the acid-base state and the adequacy of pulmonary ventilation and tissue oxygenation and includes the following signs: blood acidity level, partial pressure of carbon dioxide in arterial blood, blood bicarbonate level, and haemoglobin level. The third factor reflects the functional state of the excretory system and the state of water-electrolyte metabolism and includes the following signs: the concentration of sodium and chlorine ions and the level of blood urea.

After calculating the scores of the selected factors during the course of the traumatic disease (on the 1st, 3rd, 7th, and 14th days), a comparative analysis of the average values of factors in patients with favourable and unfavourable outcomes was carried out, which showed that the division of groups is most clearly based on the first factor.

Further analysis was aimed at studying the signs forming the first factor (LII, NLR, Mono, Lym) separately. Comparative analysis of the signs forming the first factor showed that statistically significant differences in the mean values among the patients with favourable and unfavourable outcomes simultaneously on days 1, 3, 7, and 14 were observed only in the LII sign. These results allow us to conclude that the LII is an independent objective criterion reflecting the severity of the physiological status of the patients with concomitant midface trauma and the severity of the course of the traumatic disease as a whole.

The development of a method for predicting the immediate outcomes of a concomitant midface trauma began with the search for a 'critical' time point for making a forecast. To address this task, a method of statistical analysis was applied - factor analysis (PCA) with the inclusion of a set of 22 studied physiological signs, as well as the group affiliation of the patients, depending on the outcome. This analysis was carried out in a separate form with data collected on days 1, 3, 7, and 14.

The obtained results allowed us to conclude that, taking into account the set of physiological signs included in our work, the most optimal time point (critical point) for predicting the immediate outcomes of a concomitant midface trauma is the third

day after the injury, where the sign of the group affiliation of the patients 'outcome' was included only the first factor with a factor load of 0.84.

When developing the forecasting method, the method of multivariate statistical analysis - discriminant analysis (DA) was used. The analysis included 26 signs: 22 studied physiological signs, the severity of injuries according to the military field surgery scale - Injury (MT), the age of the patients, data on the development of cerebral oedema in the patients after trauma (CE) and on the history of diseases of the cardiovascular system (CVD).

A method for predicting the immediate outcomes of a concomitant midface trauma has been developed in two variants: with and without the inclusion of signs of blood gas composition. The first variant of the discriminant function (with the inclusion of signs of blood gas composition) included the following 9 signs: age, CE, CVD, LII, blood glucose level, concentration of potassium ions in the blood, INR, pCO₂, and pO₂. The classification in the sample was 100% (without inversions).

The results of the discriminant analysis showed that the probability of a fatal outcome increases with increasing values of signs of LII, pO₂, BG, INR, pCO₂ (on day 3, respectively), age, cerebral oedema in patients after trauma (CE), a history of CVD, and a decrease in concentration of potassium ions in the blood (on day 3). At the same time, the highest factor load was observed in the sign of CE.

The second variant of the discriminant function (excluding signs of blood gas composition: SO₂, pH, pCO₂, HCO₃⁻, pO₂) included eight of the following signs: absolute number of leukocytes (on day 1), log LII (on day 3), potassium ion concentration (on day 3), log BU (on day 3), log Lym (on day 3), CE, age, and CVD. The classification in the sample was 98.4% (1 inversion).

The results of the discriminant analysis showed that the probability of a fatal outcome increases with an increase in the values of LII and BU levels (on day 3, respectively), age, cerebral oedema in patients after trauma (CE), a history of CVD, as well as a decrease in the absolute number of leukocytes (on day 1), lymphocytes (on day 3), and the concentration of potassium ions (on day 3). As in the first variant, the highest factor load was observed in the sign of CE.

A comparative analysis of the results of treatment of patients with severe concomitant midface injury showed that the complicated course of traumatic disease in the patients of the first retrospective group was almost three times more common than in the patients of the first observed group (47.6% and 14.8%, respectively). Life-threatening complications (cerebral oedema, organ or multiple organ failure, pulmonary embolism, and purulent-septic complications) were observed in 7 (33.3%) patients from the first retrospective group. In the first observed group of patients, no such complications were detected.

The incidence of infectious complications decreased from 38.1% in the first retrospective group to 11.1% in the first observed group. There was also a decrease in the severity of these complications. Generalised infectious complications prevailed in the patients of the first retrospective group, while local complications prevailed in the first observed group.

The first variant of the course of traumatic disease occurred in 96.3% of patients of the first observed group and in 66.7% of patients of the first retrospective group. The second variant - in 3.7% and 23.8% of patients, respectively. The third variant was observed in 9.5% of patients of the first retrospective group, and in the first observed group it never happened. Overall, the individuals in the first retrospective group exhibited a more severe course of traumatic disease.

The average duration of treatment of patients with severe concomitant midface trauma in ICU wards decreased from 4.9 ± 3.5 days in the first retrospective group to 3.1 ± 1 days in the first observed group ($p=0.02$). The average duration of hospitalisation also decreased from 18.3 ± 8 days in the patients of the first retrospective group to 14.2 ± 7.9 days in the first observed group ($p=0.08$).

An unfavourable outcome of the severe concomitant midface injury was observed in 4 (19%) patients of the first retrospective group. The immediate outcome of the injury turned out to be favourable for all the patients of the first observed group.

A comparative analysis of the results of treatment of patients with extremely severe concomitant midface trauma showed that the number of patients with a complicated course of traumatic disease decreased from 83.3% in the second

retrospective group to 71.4% in the first observed group. The first variant of the course of traumatic disease was observed in 33.3% of patients of the second observed group and in 19.4% of patients of the second retrospective group; the second variant of the course of traumatic disease was observed in 19% and 27.8% of patients, respectively. The third variant prevailed both in the second observed group (47.6%) and in the second retrospective group (52.8%).

The frequency of life-threatening complications (cerebral oedema, organ or multiple organ failure, pulmonary embolism, and purulent-septic complications) decreased from 80.6% in the second retrospective group to 61.9% in the second observed group.

The number of patients with a fatal outcome of an extremely severe concomitant midface injury decreased from 75% in the second retrospective group to 57.1% in the second observed group. The outcome of an extremely severe concomitant injury with a decompensated course of traumatic disease (variant 3) turned out to be unfavourable in all the patients of the second retrospective group and in 90% of patients of the second observed group. With a subcompensated course of traumatic disease (variant 2), an unfavourable outcome of combined trauma was observed in 80% of patients of the second retrospective group and in 75% of patients of the second observed group. In the compensated variant of traumatic disease course (variant 1), unfavourable outcomes were not observed among the compared groups of patients.

The difference in the average duration of treatment in the ICU wards of patients of the second observed group and the second retrospective group is insignificant. For patients of the second observed group, this indicator was 12.4 ± 3 days; for patients of the second retrospective group - 12.3 ± 1.2 days. The average duration of inpatient treatment in the patients of the second observed group exceeded that of the patients of the second retrospective group and amounted to 28.3 ± 5.5 days and 22.6 ± 3 days, respectively, which is primarily due to the large number of fatal outcomes in the second retrospective group.

FINDINGS

1. The presented algorithm of biometric analysis of features characterising the severity of the physiological status of patients with a concomitant midface trauma, utilising factor analysis of features by the time of waiting for outcomes, followed by measuring the scores of the identified factors over the course of traumatic disease, allows for simultaneous investigation of the physiological status characteristics of patients with fatal outcomes at the most critical time point (last day prior to an unfavourable outcome), as well as studying the dynamics of factor changes over the course of traumatic disease without the risk of altering the structure of the identified factors.

2. The dynamics of the average values of the leukocyte intoxication index, the neutrophil-lymphocyte ratio, and the absolute number of lymphocytes and monocytes reflects the course of the traumatic disease and the severity of the physiological status of patients with a concomitant midface trauma. The division of the cohort of patients with favourable and fatal outcomes according to these signs is the clearest on the third day after the occurrence of the injury. LII is an independent objective criterion reflecting the severity of the physiological status of the patients with a concomitant midface trauma and the severity of the course of the traumatic disease as a whole.

3. Taking into account the set of signs and time points studied by us, the third day after the occurrence of the injury is the most optimal point for predicting the immediate outcomes of the concomitant midface trauma. The methodology for predicting the immediate outcomes of a concomitant midface trauma has two variants: the first variant - with the analysis of blood gas composition, where the classification by sample was 100%; and the second variant without the analysis of blood gas composition, where the classification by sample was 98.4%.

4. In the cohort of patients with a severe concomitant midface trauma (MFS - I (MT): from 1 to 12 points)) the use of multi-stage surgical treatment tactic, taking into account the severity of the physiological status of the patients, led to a decrease in

the frequency of infectious complications by 27% (from 38.1% to 11.1%), non-infectious complications - by 5.8% (from 9.5% to 3.7%) and a decrease in mortality by 19% (from 19% to 0%); in the cohort of patients with an extremely severe concomitant midface trauma (MFS-I (MT) ≥ 12 points)) - to a decrease in the frequency of infectious complications by 9.2% (from 80.6% to 71.4%), non-infectious complications by 25.4% (from 77.8% to 52.4%) and a decrease in mortality by 17.9% (from 75% to 57.1%).

PRACTICAL RECOMMENDATIONS

1. The treatment of patients with a concomitant midface trauma should be carried out in the trauma centre of the 1st level, where the necessary physiological status for comprehensive diagnosis of patients and the implementation of an interdisciplinary approach to treatment are available.

2. Decisions regarding the amount of necessary and permissible surgical care for patients with a severe and extremely severe concomitant midface trauma are made solely taking into account the severity of the physiological status of the patients.

3. It is advisable to treat patients with a severe and extremely severe concomitant midface trauma using the tactic of multi-stage surgical treatment. In the first period of traumatic disease (the first 12 hours after the injury), the first stage of multi-stage surgical treatment is implemented, the main purpose of which is to eliminate life-threatening consequences of the injury (elimination of asphyxia and bleeding). At the first stage, a tracheostomy and temporary fixation of bone fractures by orthopaedic method and primary soft tissue plastic surgery are performed. In the second and third periods of traumatic disease (2-10 days), the second stage of multi-stage surgical treatment is implemented, the purpose of which is to achieve temporary stabilisation of the physiological status of the patient, as well as to prevent and treat infectious complications. Ventilator therapy, infusion-transfusion therapy, antibacterial therapy, rehabilitation of potential sources of infectious complications, and, in certain cases, staged fixation of fragments using minimally invasive extra-focal osteosynthesis are carried out. At this stage, radiation diagnostics (CT) of damaged midface structures is also performed. In the fourth period of traumatic disease and after the final stabilisation of the physiological status of the patient, the reconstructive-restorative (third) stage of surgical tactic is implemented in order to eliminate aesthetic and functional disorders. At that, the scope of surgical intervention at each stage, as well as the timing of its implementation, in some cases may vary depending on the course of the traumatic disease and the severity of the physiological status of the patient.

REFERENCES

1. Agadzhanyan, V. V. Organisational and Tactical Aspects of Interhospital Transportation of Critically Injured Patients in Critical Physiological status / V. V. Agadzhanyan, A. V. Shatalin, S. A. Kravtsov // Polytrauma. – 2006. – No. 1. – pp. 23–27. [In Russian]
2. Agalaryan, A. Kh. Surgical Treatment and Lethality in Patients with Abdominal Injuries in Polytrauma / A. Kh. Agalaryan // Polytrauma. – 2014. – No. 4. – pp. 24–31. [In Russian]
3. Ayvazyan, S. A. Applied Statistics. Basics of Econometrics: Textbook for students of economics. Specialised universities. In 2 volumes / S. A. Ayvazyan, V. S. Mkhitarian. – 2nd edition, revised. – Moscow: UNITI, 2001. – Vol. 1: Probability Theory and Applied Statistics. – 656 p. [In Russian]
4. Current Aspects of the Methodology of Clinical-Statistical Analysis / B. B. Bondarenko, A. G. Bart, E. A. Demchenko [et al.] // Clinical and Experimental Cardiology / edited by E. V. Shlyakhto. – St. Petersburg: AMC, 2005. – pp. 136–148. [In Russian]
5. Analysis of Hospital Mortality in Concomitant Trauma / A. N. Voloshenyuk, N. V. Zavada, S. S. Stebunov [et al.] // Emergency Medicine. – 2012. – No. 2 (2). – pp. 67–74. [In Russian]
6. Analysis of blood coagulation abnormality signs among patients with severe combined craniofacial trauma / A. N. I. Nassar, D. Yu. Madai, A. K. Iordanishvili, M. I. Idris // Stomatology for All / Int. Dental Review. – 2022. – № 4 (101). – S. 22–27. DOI: 10.35556/idr-2022-4(101)22-27. [In Russian]
7. Analysis of Complications of Severe Concomitant Cranial-Cerebral Trauma based on the Materials of a Multidisciplinary Hospital / E. A. Lebedeva, A. D. Belyaevskiy, V. A. Sarkisyan [et al.] // Medical Almanac. – 2012. – No. 1 (20). – pp. 120–122. [In Russian]
8. Analysis of the Causes of Mortality in Patients with Severe Concomitant Trauma in the Resuscitation Department of a Multidisciplinary Hospital / A. V.

- Vlasenko, O. R. Dobrushina, V. N. Yakovlev [et al.] // *General Reanimatology*. – 2009. – Vol. 5, No. 6. – pp. 31–35. [In Russian]
9. Apartsin, K. A. *Surgery of Concomitant Injuries* / K. A. Apartsin // *Siberian Medical Journal (Irkutsk)*. – 2014. – Vol. 126, No. 3. – pp. 129–133. [In Russian]
 10. Belous, I. M. *Biometric Aspects of Surgical Tactics for Comprehensive Treatment of Concomitant Midface Trauma: specialty 14.00.27 ‘Surgery’, 14.00.21 ‘Dentistry’: dis. Candidate of Medical Sciences* / Belous Inna Mikhailovna. – Veliky Novgorod, 2005. – 173 p. – Place of defence: State Educational Institution Novgorod State University named after Yaroslav the Wise. [In Russian]
 11. *Biometric Aspects of Treatment of Concomitant Trauma* / E. K. Gumanenko, Yu. A. Shcherbuk, M. G. Silyuk [et al.] // *Bulletin of Surgery named after I. I. Grekov*. – 2018. – No. 177 (3). – pp. 25–30. DOI 10.24884/0042-4625-2018-177-3-25-30. [In Russian]
 12. *Biometric Analysis of Treatment Effectiveness: Scientific edition* / V. R. Weber, D. Yu. Madai, A. G. Bart, et al. - Veliky Novgorod: Novgorod State University Press, 2005. - 150 p. [In Russian]
 13. Bogomolov, M. S. *Microsurgical Replantation of Hand Fragments* / M. S. Bogomolov, V. M. Sedov. - St. Petersburg: Elbi-SPb, 2003. - 244 p. [In Russian]
 14. Boymuradov, Sh. A. *The Value of Computed Tomography in Patients with Fractures of the Zygomatic Bone and Zygomatic Arch* / Sh. A. Boymuradov // *Russian Otorhinolaryngology*. - 2009. - No. 4 (41). - pp. 38-43. [In Russian]
 15. Bondarenko, A. V. *Hospital Mortality in Concomitant Trauma and Possibilities of its Reduction* / A. V. Bondarenko, V. A. Peleganchuk, O. A. Gerasimova // *Herald of Traumatology and Orthopedics named after N. N. Priorov*. - 2004. - No. 3. - p. 49. [In Russian]
 16. Vinogradova, N. G. *Optimisation of Surgical Tactics in Patients with Midface Trauma: specialty 14.01.15 ‘Traumatology and Orthopedics’: thesis for the degree of Candidate of Medical Sciences* / Vinogradova Natalya Gennadievna. - Perm, 2015. - 165 p. – Place of defence: State Educational Institution Perm

- State Medical University named after Academician E. A. Wagner of the Ministry of Healthcare of the Russian Federation. [In Russian]
17. Vlasov, A. M. Diagnosis and Treatment of Concomitant Cranial-Cerebral and Maxillofacial Trauma: specialty 14.00.21 'Dentistry', 14.00.28 'Neurosurgery': author's abstract of the thesis for the degree of Candidate of Medical Sciences / Vlasov Andrey Mikhailovich; Omsk State Medical Academy. - Omsk, 2003. - 15 p. [In Russian]
 18. Vliyanie ostroi posttravmaticheskoi koagulopatii na blizhaishie iskhody tyazheloi sochetannoi cherepno-litsevoi travmy / D. Yu. Madai, A. K. Iordanishvili, M. I. Idris, K. A. Absava, A. N. I. Nassar // Sovremennye podkhody k diagnostike i lecheniyu khirurgicheskikh zabolevanii, travm i ikh oslozhnenii : Sb. tr. Obscheros. mezhvedomstvennoi nauch.-prakt. konf. / Departament med. obespecheniya Federal'noi sluzhby voisk natsional'noi gvardii Ros. Federatsii, FGKUZ Glavnyi voennyi klinicheskii gospital' voisk natsional'noi gvardii Ros. Federatsii, FGBOU DPO RMANPO Minzdrava Rossii, Med. in-t nepreryvnogo obrazovaniya FGBOU VO MGUPP ; [red. kol.: A. N. Ivashchenko, D. R. Ivchenko, V. F. Zubritskii [i dr.]. – M. : Red. zhurnala «Na boevom postu», 2022. – S. 143–144. [In Russian]
 19. Vliyanie soputstvuyushchikh zabolevanii na blizhaishie iskhody sochetannoi cherepno-litsevoi travmy / A. N. I. Nassar, M. I. Idris, D. Yu. Madai, A. K. Iordanishvili // Aktual'nye voprosy chelyustno-litsevoi khirurgii i stomatologii : Materialy. Vseros. nauch.-prakt. konf., posvyashch. 100-letiyu so dnya rozhdeniya professora V. A. Malysheva (Sankt-Peterburg, 23-24 noyabrya 2022 g.) / Glavnoe voenno-meditsinskoe upravlenie M-va oborony Ros. Federatsii, Voенno-meditsinskaya akademiya im. S. M. Kirova, Assotsiatsiya obshchestvennykh ob"edinenii «Stomatologicheskaya Assotsiatsiya Rossii» [i dr.]. – SPb. : Izd-vo Voен.-med. akad. im. S. M. Kirova, 2022. – S. 186–188. [In Russian]

20. Military Field Surgery: textbook / edited by E. K. Gumanenko. - 2nd edition, revised and supplemented. - Moscow: GEOTAR-Media, 2022. - 768 p. [In Russian]
21. Voloshenyuk, A. N. Socio-Economic Significance of Polytrauma / A. N. Voloshenyuk, S. V. Filinov // Military Medicine. - 2011. - No. 1 (18). - pp. 118-120. [In Russian]
22. Gabdulkhakov, R. M. Prognosis and Intensive Therapy in Concomitant Trauma: specialty 14.00.37 'Anesthesiology and Resuscitation', 14.00.27 'Surgery': author's abstract of the thesis for the degree of Doctor of Medical Sciences / Gabdulkhakov Rail Munirovich. - Moscow, 2009. - 47 p. – Place of defence: Russian State Medical University. [In Russian]
23. Gerasimov, L. V. Selected Issues of Pathogenesis and Intensive Treatment of Severe Concomitant Trauma / L. V. Gerasimov, N. A. Karpun, O. S. Pirozhkova // General Reanimatology. - 2012. - Vol. 8, No. 4. - pp. 111-117. [In Russian]
24. Golovko, K. P. Implementation of a Multi-Stage Surgical Treatment Method in Patients with Severe Concomitant Injuries of the Maxillofacial Region / K. P. Golovko, D. Yu. Madai // Health. Medical Ecology. Science. - 2012. - No. 1-2 (47-48). - pp. 38-39. [In Russian]
25. Golovko, K. P. Modern Approach to Comprehensive Treatment of Concomitant Injuries of the Maxillofacial Region: specialty 14.01.17 'Surgery', 14.01.14 'Dentistry': thesis for the degree of Doctor of Medical Sciences / Golovko Konstantin Petrovich. - St. Petersburg, 2015. - 331 p. – Place of defence: Federal State Budgetary Educational Institution Military Medical Academy named after S. M. Kirov of the Ministry of Defence of the Russian Federation. [In Russian]
26. Gumanenko, E. K. Concomitant Injuries from the Perspective of Objective Assessment of Trauma Severity: specialty 14.00.27 'Surgery': author's abstract of the thesis for the degree of Doctor of Medical Sciences / Gumanenko Evgeny Konstantinovich; Military Medical Academy named after S. M. Kirov of the Ministry of Defence of the Russian Federation. - St. Petersburg, 1992. - 50 p. [In Russian]

27. Guseynov, A. G. Optimisation of Treatment for Concomitant Trauma in Road Traffic Accidents / A. G. Guseynov // International Journal of Applied and Fundamental Research. - 2011. - No. 9. - pp. 92-93. [In Russian]
28. Danilevich, M. O. Severe Craniofacial Trauma: Features of Clinical Course and Multidisciplinary Approach to Comprehensive Treatment: specialty 14.01.18 'Neurosurgery': thesis for the degree of Doctor of Medical Sciences / Danilevich Marina Olegovna; Russian Scientific Research Neurosurgical Institute named after A. L. Polenov. - St. Petersburg, 2016. - 283 p. – Place of defence: North-Western Federal Medical Research Centre named after V. A. Almazov of the Ministry of Healthcare of the Russian Federation. [In Russian]
29. Epifanov, S. A. Computer Technologies in Reconstructive Surgery of the Midface / S. A. Epifanov, V. N. Balin, S. S. Khrykov // Bulletin of the National Medical and Surgical Centre named after N. I. Pirogov. - 2014. - Vol. 9, No. 4. - pp. 28-31. [In Russian]
30. Ermolov, A. S. Choice of Criteria for Assessing the Severity of Closed Concomitant Trauma / A. S. Ermolov, V. I. Kartavenko, M. M. Abakumov // Assistance in Concomitant Trauma: Collection of scientific works. - Moscow, 1997. - pp. 11-14. [In Russian]
31. Inozemtsev, E. O. Current Issues in the Surgery of Concomitant Injuries (based on publications in the journal 'Politravma') / E. O. Inozemtsev, E. G. Grigoryev, K. A. Apartsin // Politravma. - 2017. - No. 1. - pp. 6-11. [In Russian]
32. Integrated Systems for Assessing the Severity of Patients' Physiological status in Polytrauma / B. R. Gelfand, A. I. Yaroshevsky, D. N. Protsenko, Yu. Ya. Romanovsky // Bulletin of Intensive Therapy. - 2004. - No. 1. - pp. 58-65. [In Russian]
33. Integration of Severity Criteria for Polytrauma with the International Classification of Diseases / V. V. Agadzhanian, S. A. Kravtsov, I. A. Zheleznikova, et al. // Politravma. - 2014. - No. 1. - pp. 6-14. [In Russian]
34. Kartavenko, V. I. Application of the APACHE-II Scale for Assessing the Severity of Patient Injuries / V. I. Kartavenko, A. K. Shabanov // Integral

- Assessment and Forecasting in Emergency Medicine: Proceedings of the Scientific and Practical Conference Dedicated to the 70th Anniversary of Prof. Yu. N. Tsibin (St. Petersburg, November 9-10, 2000): Collection of Works / Committee on Health of the Administration of St. Petersburg, St. Petersburg Scientific Research Institute of Emergency Medical Care named after I. I. Dzhanelyidze; edited by S. F. Bagnenko. - St. Petersburg, 2001. - pp. 18-26. [In Russian]
35. Clinical and biostatistical approach to assessing the severity of physical status of patients with combined midface trauma. Part 1 / A. N. I. Nassar, V. A. Bart, D. Yu. Madai, A. K. Iordanishvili, M. I. Idris // *Stomatology for All / Int. Dental Review*. – 2023. – № 1 (102). – S. 10-17 DOI: 10.35556/idr-2023-1(102)10-16. [In Russian]
36. Clinical and biostatistical approach to assessing the severity of physical status of patients with combined midface trauma. Part 2/ A. N. I. Nassar, V. A. Bart, D. Yu. Madai, A. K. Iordanishvili, M. I. Idris // *Stomatology for All / Int. Dental Review*. – 2023. – № 2 (103). – S. 32-38. DOI: 10.35556/idr-2023-2(103)32-38. [In Russian]
37. Klypa, T. V. Hyperglycemia in Critical Physiological status / T. V. Klypa, M. S. Orekhova, L. I. Zabrosaeva // *Diabetes Mellitus*. - 2015. - Vol. 18, No. 1. - pp. 33-41. DOI 10.14341/DM2015133-41. [In Russian]
38. Kolesnikov, A. N. Algorithms for Assessing the Severity and Shock Therapy in Patients with Polytrauma during Peacetime and Wartime (literature review) / A. N. Kolesnikov, A. M. Pliev, O. S. Antropova // *Modern Medicine: New Approaches and Current Research: Collection of Materials of the International Scientific and Practical Conference Dedicated to the 80th Anniversary of Chechen State University (Grozny, September 26-27, 2018) / Federal State Budgetary Educational Institution Chechen State University*; edited by M. R. Nahaev. - Grozny: Publishing House of Chechen State University, 2018. - pp. 347-366. [In Russian]

39. Comprehensive Approach to Evaluating Experimental Studies / D. Yu. Madai, V. R. Weber, A. S. Grivkov, et al. // International Journal of Experimental Education. - 2013. - No. 11-2. - pp. 40-46. [In Russian]
40. Craniofacial Trauma / S. A. Eolchiyan, A. A. Potapov, F. A. Van Damm, et al. // Clinical Guide to Cranial Trauma: in 3 volumes / edited by A. N. Konovalov, L. B. Likheterman, A. A. Potapov; N. N. Burdenko Institute of Neurosurgery, RAMS. - Moscow: ANTIDOR, 2002. - Vol. 3: Consequences and Complications of Cranial Trauma, Standards and Recommendations, Neurorehabilitation, Expertise. - Chap. 16. - pp. 313-346. [In Russian]
41. Criteria for Assessing the Physiological status of Patients with Polytrauma during Interhospital Transportation / V. V. Agadzhanyan, A. V. Shatalin, S. A. Kravtsov, D. A. Skopintsev // Politravma. - 2011. - No. 1. - pp. 5-11. [In Russian]
42. Lang, T. A. How to Describe Statistics in Medicine: A Guide for Authors, Editors, and Reviewers / T. A. Lang, M. Sesik; translated from English, edited by V. P. Leonov. - Moscow: Practical Medicine, 2011. - 480 p. [In Russian]
43. Larin, M. A. Analysis of the Causes of Mortality in Severe Concomitant Trauma / M. A. Larin, V. P. Ryazantsev // Bulletin of the Eastern Siberian Scientific Centre of the Siberian Branch of the Russian Academy of Medical Sciences. - 2005. - No. 3 (41). - pp. 110-111. [In Russian]
44. Lebedev, V. V. Emergency Neurosurgery: A Guide for Physicians / V. V. Lebedev, V. V. Krylov. - Moscow: Medicine, 2000. - 568 p. [In Russian]
45. Madai, D. Yu. Analiz svyazei pokazatelei klinicheskogo analiza krovi i tyazhesti povrezhdenii u postradavshikh s tyazheloi sochetannoi cherepno-litsevoi travmoi / D. Yu. Madai, M. I. Idris, A. N. I. Nassar // Nauka SPbGU – 2020 : Sbornik materialov Vseros. konf. po estestvennym i gumanitarnym naukam s mezhdunar. uchastiem (Sankt-Peterburg, 24 dekabrya 2020 g.). – SPb. : Skifiya-print, 2021. – S. 701–702. [In Russian]
46. Madai, D. Yu. Comprehensive Use of Polysaccharides for Correction of Post-Aggressive Adaptive Reactions (PAR) in Patients with Inflammatory Diseases and Injuries of the Maxillofacial Region (Theoretical aspects of the problem,

- experimental and clinical research): specialty 14.00.21 'Dentistry': thesis for the degree of Doctor of Medical Sciences / Madai Dmitry Yurievich; Military Medical Academy named after S. M. Kirov of the Ministry of Defense of the Russian Federation. - St. Petersburg, 1998. - 44 p. [In Russian]
47. Madai, D. Yu. Treatment of Fractures of the Nasoethmoidal Complex in Patients with Severe Craniofacial Trauma: Clinical-Functional Substantiation of Tactics / D. Yu. Madai, E. L. Sokirko, A. Yu. Shcherbuk; St Petersburg University - St Petersburg: Publishing House of St Petersburg University, 2014 - 109 p. - (Surgery). [In Russian]
48. Madai, D. Yu. Oslozhneniya i prichiny letal'nogo iskhoda u postradavshikh s neblagopriyatnym iskhodom tyazheloi sochetannoi cherepno-litsevoi travmy / D. Yu. Madai, M. I. Idris, A. N. I. Nassar // Nauka SPbGU – 2020 : Sbornik materialov Vseros. konf. po estestvennym i gumanitarnym naukam s mezhdunar. uchastiem (Sankt-Peterburg, 24 dekabrya 2020 g.). – SPb. : Skifiya-print, 2021. – S. 703–704. [In Russian]
49. Madai, D. Yu. Concomitant Craniofacial Trauma / D. Yu. Madai; Ministry of Education and Science of the Russian Federation, Yaroslav-the-Wise Novgorod State University. - Veliky Novgorod: Publishing House of NovSU named after Yaroslav-the-Wise, 2011 - 175 p. [In Russian]
50. Madai, D. Yu. Terminology and Classification of Combat and Non-Combat Surgical Trauma: a textbook / D. Yu. Madai, E. K. Gumanenko, I. M. Samokhvalov; St Petersburg University. - 2nd edition, revised - St Petersburg: Publishing House of St Petersburg University, 2013 - 55 p. - (Surgery). [In Russian]
51. Madai, O. D. Substantiation of Programmed Multi-Stage Surgical Treatment of Injuries to the Midface Region in Patients with Concomitant Trauma: specialty 14.03.01 'Human Anatomy', 14.01.17 'Surgery': thesis for the degree of Candidate of Medical Sciences / Madai Olga Dmitrievna. - St. Petersburg, 2019 - 177 p. - Place of defence: S. M. Kirov Military Medical Academy of the Ministry of Defence of the Russian Federation. [In Russian]

52. Malanin, D. A. Methods of Objective Assessment of Trauma Severity and their Practical Application: methodological recommendations / D. A. Malanin, O. Yu. Bosko. - Volgograd: Publishing House of Volgograd State Medical University, 2008 - 15 p. [In Russian]
53. Malanchuk, V. A. Possibilities of Orbit and Adnexal Apparatus Reconstruction in Patients with Midface Injuries / V. A. Malanchuk, E. A. Astapenko, Yu. V. Chepurnoy // Modern dentistry. - 2013. - No. 2 (57). - P. 46-48. [In Russian]
54. Medvedev, Yu. A. Combined Injuries of the Midface Zone. Part I / Yu. A. Medvedev, P. S. Petruk // Head and Neck/Russian Edition. Journal of the All-Russian Public Organisation Federation of Specialists in the Treatment of Head and Neck Diseases. - 2017. - No. 1. - P. 51-55. [In Russian]
55. Mediko-epidemiologicheskie aspekty travmaticheskikh povrezhdenii srednei zony litsa / D. Yu. Madai, A. K. Iordanishvili, M. I. Idris, A. N. I. Nassar // Meditsina i obrazovanie. – 2022. – № 2 (10). – S. 12–16. [In Russian]
56. Methodological Aspects of Surgical Tactics in Patients with Severe Concomitant Craniofacial Trauma Considering the Severity of Traumatic Disease / Yu. A. Shcherbuk, D. Yu. Madai, S. V. Gavrilin [et al.] // Bulletin of Surgery named after I. I. Grekov. - 2014. - Vol. 173, No. 3. - P. 49-54. [In Russian]
57. Mirenkov, K. B. Treatment of Concomitant Injuries and Diseases of the Limbs / K. V. Mirenkov, T. A. Amro, S. A. Melashenko // Materials of All-Russian Scientific Conference. - Moscow, 2003. - P. 229-230. [In Russian]
58. Multi-Stage Surgical Treatment for Reducing Lethality in Patients with Concomitant Maxillofacial and Cranio-Cerebral Trauma / D. Yu. Madai, K. P. Golovko, V. I. Badalov [et al.] // Emergency Medical Care. - 2016. - Vol. 17, No. 2. - P. 33-41. [In Russian]
59. Novozhilov, A.V., Kosenkova, D.V., Zaitsev, A.P., Aparcin, K.A. Monitoring of Trauma in Concomitant Mechanical Injuries. Politravma. 2007; (1): 9-14. [In Russian]

60. Nabiev, E.N., Tezekbaev, K.M., Tusupov, D.M. Methods of Objective Assessment of Injury Severity in Polytrauma. Bulletin of the Kyrgyz State Medical Academy named after I.K. Akhunbaev. 2018; (5-6): 80-88. [In Russian]
61. Nassar, A. N. I. Obosnovannaya khirurgicheskaya taktika v kompleksnom lechenii postradavshikh s sochetannymi povrezhdeniyami srednei zony litsa / A. N. I. Nassar, D. Yu. Madai // Meditsinskaya pomoshch' pri travmakh. Novoe v organizatsii i tekhnologiyakh. Perspektivy importozameshcheniya v Rossii : Pyaty yubileinyi kongress s mezhdunar. uchastiem (Sankt-Peterburg, 28–29 fevralya 2020 g.) : sb. tez. / M-vo zdravookhraneniya Ros. Federatsii, Otd-nie med. nauk RAN, Komitet po zdravookhraneniyu Sankt-Peterburga [i dr.]. – SPb. : Chelovek i ego zdorov'e, 2020. – S. 166–167. [In Russian]
62. Nassar, A. N. I. Scoring Models For The Severity Of Combined Craniofacial Trauma (A Review) / A. N. I. Nassar, D. Yu. Madai // Kubanskii nauchnyi meditsinskii vestnik. – 2020. – T. 27, № 5. – S. 144–162. [In Russian]
63. Nikolaeva, E.V. Trauma of Extremities: Medical Expertise. Moscow: Grant, 2003. 239 p. [In Russian]
64. Trauma severity scoring — a reserve for a favorable treatment outcome of patients with craniofacial trauma/ A. N. I. Nassar, M. I. Idris, D. Yu. Madai, K. A. Absava // Stomatology for All / Int. Dental Review. – 2020. – № 1 (90). – S. 16–23. DOI: 10.35556/idr-2020-1(90)16-22. [In Russian]
65. Gumanenko, E.K., Boyarintsev, V.V., Suprun, T.Yu., Lyashedko, P.P. Objective Assessment of Injury Severity: textbook. St Petersburg: Military Medical Academy named after S.M. Kirov, 1999. 110 p. [In Russian]
66. Sokolov, V.A., Byalik, E.I., Takiev, A.T., Boyarshinova, O.I. Surgical Treatment of Fractures of the Distal Femur in Patients with Concomitant and Multiple Trauma. Bulletin of Traumatology and Orthopaedics named after N.N. Priorov. 2004; (1): 20-26. [In Russian]
67. Berezka, N.I., Litovchenko, V.A., Garyachiy, E.V., et al. Optimisation of Surgical Treatment Tactics for Patients with Polytrauma Using Severity

- Assessment Scales. Scientific Bulletin of the Belgorod State University. Series: Medicine. Pharmacy. 2014; (4, 175): 116-119. [In Russian]
68. Sorokin, E.P., Gritsan, A.I., Ponomarev, S.V., Shilyaeva, E.V. Experience of Using Prognostic Scales to Assess Survivability in Patients with Concomitant Chest and Abdominal Injuries. Bulletin of Anaesthesiology and Intensive Care. 2013; 10(5): 47-50. [In Russian]
 69. Tulupov, A.N., Besaev, G.M., Sinenchenko, G.I., et al. Assessment of Injury Severity and Dynamic Forecasting in Concomitant Trauma: educational-methodical guide. St Petersburg: Stix, 2014. 53 p. [In Russian]
 70. Patent No. 105151 Russian Federation, IPC A61B 17/00 (2006.01). Method of non-focal-stem osteosynthesis in Le Fort-type fractures of the upper jaw - HEGEN: No. 2009148550: filed on 22 December 2009: published on 10 June 2011. Belevitin, A.B., Golovko, K.P., Madai, D.Yu., Samokhvalov, I.M.; patent holder Golovko, K.P. - 15 p. [In Russian]
 71. Patent No. 2430698 Russian Federation, IPC A61B 17/60 (2006.01). Method of non-focal-stem osteosynthesis in Le Fort-type fractures of the upper jaw - II medium and a set for its implementation: No. 2009148548: filed on 22 December 2009: published on 10 October 2011. Belevitin, A.B., Golovko, K.P., Madai, D.Yu., Samokhvalov, I.M.; patent holder Golovko, K.P. - 9 p. [In Russian]
 72. Grishenkova, L.N., Gorbachev, F.A., Anishchenko, S.L., et al. Fractures of the Bones of the Midfacial Region: Forensic Medical Examination. Textbook. Minsk: Vysh. Shk., 2015. 177 p. [In Russian]
 73. Belevitin, A.B., Parfenov, V.E., Samokhvalov, I.M., et al. Prospects for the Use of Endovideosurgery in the Treatment of Upper and Midface Injuries. Bulletin of the Russian Military Medical Academy. 2009; (4, 28): 96-103. [In Russian]
 74. Stukanov, M.M., Yudakova, T.N., Maximishin, S.V., et al. Indicators Associated with Fatal Outcomes in Patients with Traumatic Shock. Politravma. 2015; (2): 37-43. [In Russian]

75. Humanenko, E.K., Kozlov, V.K. Polytrauma: Traumatic Disease, Immune System Dysfunction, and Modern Treatment Strategy. Edited by E.K. Humanenko, V.K. Kozlov. Moscow: GEOTAR-Media, 2008. 608 p. [In Russian]
76. Puras, Yu.V., Talybov, A.E., Krylov, V.V. Treatment Outcomes in Patients with Concomitant Cranial-Brain Injury at the In-patient Stage of Medical Care. Emergency Medical Care. Journal named after N.V. Sklifosovsky. 2013; (1): 38-45. [In Russian]
77. Pushkov, A.A. Concomitant Trauma: a View of a Practical Surgeon. Rostov State Medical University, Emergency Hospital No. 2. Rostov-on-Don: Feniks, 1998. 315 p. (Hippocrates). [In Russian]
78. Rakhmanov, A.T., Yuldashev, I.M., Lesogorov, S.F., et al. Prevalence of Midface Injuries. Bulletin of the Kyrgyz-Russian Slavic University. 2018; (6, 18): 75-77. [In Russian]
79. Seliverstov, P.A., Shapkin, Yu.G. Assessment of Severity and Outcome Prediction in Polytrauma: Current State of the Issue (review). Modern Technologies in Medicine. 2017; 9(2): 207-218. [In Russian]
80. Semenov, A.V., Novikova, O.V. Features of Treatment of Cranial-Brain Trauma Combined with Maxillofacial Injuries in a Neurosurgical Hospital. Bulletin of the Eastern Siberian Scientific Centre, Siberian Branch of the Russian Academy of Medical Sciences. 2005; (3, 41): 138-139. [In Russian]
81. Semenov, A.V., Sorokovikov, V.A. Scales for Assessing Severity and Outcome Prediction in Trauma. Politravma. 2016; (2): 80-90. [In Russian]
82. Modern Concepts of Diagnosis and Reconstructive-Restorative Treatment of Patients with Post-Traumatic Defects and Deformities of the Facial Skeleton. Drobyshev, A.Yu., Kobzeva, I.V., Dubina, L.Kh., et al. Bulletin of Experimental and Clinical Surgery. 2012; 5(1): 181-185. [In Russian]
83. Sokolov, V.A., Klopov, L.G. Clinical and Statistical Classification of Concomitant and Multiple Injuries with Elements of Prediction. Orthopaedics, Traumatology, and Prosthetics. 1990; (10): 54-57. [In Russian]

84. Sokolov, V.A. Multiple and Concomitant Injuries: (practical guide for trauma doctors). Moscow: GEOTAR-Media, 2006. 512 p. [In Russian]
85. Solodovnikov, P.N. Treatment of Concomitant Injuries and Diseases of the Limbs. Materials of the All-Russian Scientific Conference. Moscow, 2003. 293-294. [In Russian]
86. Mironov, S.P., Kakorina, E.P., Andreeva, T.M., Ogryzko, E.V. State of Trauma and Orthopaedic Care for the Population of the Russian Federation. Bulletin of Traumatology and Orthopaedics named after N.N. Priorov. 2007; (3): 3-10. [In Russian]
87. Combined Mechanical Trauma: Guide for Physicians. Edited by A.N. Tulupov. St. Petersburg: Stiks, 2012. 395 p. [In Russian]
88. Madai, D.Yu., Shcherbuk, A.Yu., Absava, K.A., et al. Strategy for Specialised Medical Care for Patients with Craniofacial Trauma in a Level 1 Trauma Centre. Aspirant Doctor. 2013; 61(6.1): 126-132. [In Russian]
89. Tactics of Programmed Multi-Stage Surgical Treatment in Patients with Severe Cranio-Facial Trauma. Madai, D.Yu., Golovko, K.P., Badalov, V.I., et al. Emergency Medical Care - 2015: Abstracts of All-Russian Scientific and Practical Conference (St Petersburg, June 25-26, 2015). St Petersburg: Publishing House of St Petersburg Medical University, 2015. 79-80. [In Russian]
90. Timofeev, A.A. Manual on Maxillofacial Surgery and Surgical Dentistry. Kiev: Chervona Ruta-Ture, 2002. 1024 p. [In Russian]
91. Traumatology: National Guidelines. Quality Medical Society Association; edited by G.P. Kotelnikov, S.P. Mironov. 2nd edition, revised and supplemented. Moscow: GEOTAR-Media, 2011. 1100 p. (national guidelines). [In Russian]
92. Tulupov, A.N. Severe Concomitant Trauma. Edited by B.I. Djurko. St Petersburg: RA 'Russkiy Yuvilyr', 2015. 314 p. (Library of Emergency Care Doctors). [In Russian]
93. Khaanin, M.Yu. Correction of Anaemia in Multiple and Concomitant Injuries of the Musculoskeletal System. Medical Bulletin of Bashkortostan. 2012; 7(2): 78-80. [In Russian]

94. Surgical Treatment of Cranio-Orbital Injuries in the Acute Period of Cranio-Cerebral Trauma. Krylov, V.V., Levchenko, O.V., Shalumov, A.Z., Kutrovskaya, N.Yu. Paediatric Neurosurgery and Neurology. 2012; (2-3, 32-33): 119-129. [In Russian]
95. Tsibin, Yu.N. Multifactorial Assessment of the Severity of Traumatic Shock in Clinics. Grekov Bulletin of Surgery. 1980; 125(9): 62-67. [In Russian]
96. Tsibin, Yu.N., Galtseva, I.V., Rybakov, I.R. Prediction of Outcomes in Severe Trauma Complicated by Shock. Traumatic Shock: Proceedings. Leningrad: Sklifosovsky Research Institute of Emergency Medicine; edited by M.P. Gvozdev. Leningrad, 1976. 59-62. [In Russian]
97. Shabanov, A.K. The Use of Objective Methods for Assessing the Severity of Injuries and the Physiological status of Patients with Concomitant Trauma. [Candidate's thesis]. Moscow, 2006. 120 p. (Place of defence: Sklifosovsky Research Institute of Emergency Medicine). [In Russian]
98. Shchedrenok, V.V., Yakovenko, I.V., Moguchaya, O.V., et al. Clinical and Organisational Aspects of Concomitant Cranio-Cerebral Trauma. Edited by V.V. Shchedrenok. St Petersburg: Publishing House of the Russian Scientific Centre for Surgery, 2010. 434 p. [In Russian]
99. Emergency Reconstructive Operations in Craniofacial Trauma. Okonskaya, Yu.I., Totorculov, R.I., Pavlov, V.A., et al. Institute of Dentistry. 2012; (4, 57): 32-35. [In Russian]
100. Endoscopic Technologies in Reconstructive and Restorative Craniofacial Surgery: textbook for higher and continuous medical education. Madai, O.D., Gaivoronsky, I.V., Badalov, V.I., et al. St Petersburg University, Department of Maxillofacial Surgery and Surgical Dentistry, Military Medical Academy named after S.M. Kirov, Department of Normal Anatomy, Department of Military Field Surgery. St. Petersburg: Mediapapir, 2017. 50 p. [In Russian]
101. A comparison of the abilities of nine scoring algorithms in predicting mortality / J. W. Meredith, G. Evans, P. D. Kilgo [et al.] // The Journal of trauma. – 2002. – Vol. 53, № 4. – P. 621–628. DOI: 10.1097/00005373-200210000-00001.

102. A comparison of the injury severity score and the trauma mortality prediction model / A. Cook, J. Weddle, S. Baker [et al.] // The journal of trauma and acute care surgery. – 2014. – Vol. 76, № 1. – P. 47–52. DOI: 10.1097/TA.0b013e3182ab0d5d.
103. A new characterization of injury severity / H. R. Champion, W. S. Copes, W. J. Sacco [et al.] // The Journal of trauma. – 1990. – Vol. 30, № 5. – P. 539–545. DOI: 10.1097/00005373-199005000-00003.
104. A revision of the Trauma Score / H. R. Champion, W. J. Sacco, W. S. Copes [et al.] // The Journal of trauma. – 1989. – Vol. 29, № 5. – P. 623–629. DOI: 10.1097/00005373-198905000-00017.
105. A trauma mortality prediction model based on the anatomic injury scale / T. Osler, L. Glance, J. S. Buzas [et al.] // Annals of surgery. – 2008. – Vol. 247, № 6. – P. 1041–1048. DOI: 10.1097/SLA.0b013e31816ffb3f.
106. Abbreviated Injury Scale. – Text : electronic // Association for the Advancement of Automotive Medicine : website. – URL: <https://www.aaam.org/abbreviated-injury-scale-ais/> (date of request: 06.04.2020).
107. Abosadegh, M. M. Association of traumatic head injuries and maxillofacial fractures: A retrospective study / M. M. Abosadegh, S. A. Rahman, N. Saddki // Dental traumatology. – 2017. – Vol. 33, № 5. – P. 369–374.
108. Acute physiology and chronic health evaluation II and simplified acute physiology score II in predicting hospital mortality of neurosurgical intensive care unit patients / S. K. Park, H. J. Chun, D. W. Kim [et al.] // Journal of Korean medical science. – 2009. – Vol. 24, № 3. – P. 420–426. DOI: 10.3346/jkms.2009.24.3.420.
109. Admission heart rate is a predictor of mortality / E. J. Ley, M. B. Singer, M. A. Clond [et al.] // The journal of trauma and acute care surgery. – 2012. – Vol. 72, № 4. – P. 943–947. DOI: 10.1097/TA.0b013e3182465527.
110. An international normalized ratio-based definition of acute traumatic coagulopathy is associated with mortality, venous thromboembolism, and multiple organ failure after injury / I. D. Peltan, L. K. Vande Vusse, R. V. Maier,

- T. R. Watkins // *Critical care medicine*. – 2015. – Vol. 43, № 7. – P. 1429–1438. DOI: 10.1097/CCM.0000000000000981.
111. APACHE II: a severity of disease classification system / W. A. Knaus, E. A. Draper, D. P. Wagner, J. E. Zimmerman // *Critical care medicine*. – 1985. – Vol. 13, № 10. – P. 818–829.
112. Assessment of injury severity: the triage index / H. R. Champion, W. S. Copes, D. S. Hannan [et al.] // *Critical care medicine*. – 1980. – Vol. 8, № 4. – P. 201–208. DOI: 10.1097/00003246-198004000-00001.
113. Blood glucose concentration does not affect outcome in brain trauma: A 31P MRS study / R. Vink, E. M. Golding, J. P. Williams, T. K. McIntosh // *Journal of cerebral blood flow and metabolism*. – 1997. – Vol. 17, № 1. – P. 50–53. DOI: 10.1097/00004647-199701000-00007.
114. Blood glucose concentrations in prehospital trauma patients with traumatic shock: A retrospective analysis / J. Kreutziger, W. Lederer, S. Schmid [et al.] // *European journal of anaesthesiology*. – 2018. – Vol. 35, № 1. – P. 33–42. DOI: 10.1097/EJA.0000000000000733.
115. Blood glucose levels may aid the decision for CT scan in minor head trauma / G. A. Alexiou, A. Sotiropoulos, G. D. Lianos [et al.] // *Disease markers*. – 2019. – Vol. 2019, Article 1065254. DOI: 10.1155/2019/1065254.
116. Bosse, M. J. CAQ: Orthopaedic trauma ‘Damage control’ / M. J. Bosse // *Journal of orthopaedic trauma*. – 2007. – Vol. 21, № 1. – P. 1–4. DOI: 10.1097/BOT.0b013e318030dfd2.
117. Boyd, C. R. Evaluating trauma care: the TRISS method. Trauma Score and the Injury Severity Score / C. R. Boyd, M. A. Tolson, W. S. Copes // *The Journal of trauma*. – 1987. – Vol. 27, № 4. – P. 370–378.
118. Can serum glucose level in early admission predict outcome in patients with severe head trauma? / F. Salehpour, A. M. Bazzazi, J. Aghazadeh [et al.] // *World neurosurgery*. – 2016. – Vol. 87. – P. 132–135. DOI: 10.1016/j.wneu.2015.11.048.

119. Changes in the outcomes of severe trauma patients from 15-year experience in a Western European trauma ICU of Emilia Romagna region (1996-2010). A population cross-sectional survey study / S. Di Saverio, G. Gambale, F. Coccolini [et al.] // *Langenbeck's archives of surgery*. – 2014. – Vol. 399, № 1. – P. 109–126. DOI: 10.1007/s00423-013-1143-9.
120. Changes of immunoregulatory cells induced by psychological and physical stress: relationship to plasma catecholamines / R. M. Landmann, F. B. Müller, C. Perini [et al.] // *Clinical and experimental immunology*. – 1984. – Vol. 58, № 1. – P. 127–135.
121. Characterization of acute coagulopathy and sexual dimorphism after injury: females and coagulopathy just do not mix / J. B. Brown, M. J. Cohen, J. P. Minei [et al.] // *The journal of trauma and acute care surgery*. – 2012. – Vol. 73, № 6. – P. 1395–1400. DOI: 10.1097/TA.0b013e31825b9f05.
122. Chronic physiological status and outcomes of pediatric trauma patients / C. M. Miller, J. Shi, K. K. Wheeler [et al.] // *The journal of trauma and acute care surgery*. – 2013. – Vol. 75, № 2. – P. 250–257. DOI: 10.1097/TA.0b013e3182930fb7.
123. Classification of the severity of injury / H. J. Oestern, H. Tscherne, J. Sturm, M. Nerlich // *Der Unfallchirurg*. – 1985. – Vol. 88, № 11. – P. 465–472.
124. Comparison between the prognostic value of the white blood cell differential count and morphological parameters of neutrophils and lymphocytes in severely injured patients for 7-day in-hospital mortality / S. W. Lam, L. P. H. Leenen, W. W. van Solinge [et al.] // *Biomarkers*. – 2012. – Vol. 17, № 7. – P. 642–647. DOI: 10.3109/1354750X.2012.712161.
125. Comparison of alternative methods for assessing injury severity based on anatomic descriptors / W. J. Sacco, E. J. MacKenzie, H. R. Champion [et al.] // *The Journal of trauma*. – 1999. – Vol. 47, № 3. – P. 441–446. DOI: 10.1097/00005373-199909000-00001.
126. Comparison of APACHE II, TRISS, and a proposed 24-hour ICU point system for prediction of outcome in ICU trauma patients / M. J. Vassar, C. L. Wilkerson,

- P. J. Duran [et al.] // The Journal of trauma. – 1992. – Vol. 32, № 4. – P. 490–499. DOI: 10.1097/00005373-199204000-00014.
127. Comparison of current injury scales for survival chance estimation: an evaluation comparing the predictive performance of the ISS, NISS, and AP scores in a Dutch local trauma registration / S. P. G. Frankema, E. W. Steyerberg, M. J. R. Edwards, A. B. van Vugt // The Journal of trauma. – 2005. – Vol. 58, № 3. – P. 596–604. DOI: 10.1097/01.ta.0000152551.39400.6f.
128. Comparison of postinjury multiple-organ failure scoring systems: Denver versus Sequential Organ Failure Assessment / D. C. Dewar, A. White, J. Attia [et al.] // The journal of trauma and acute care surgery. – 2014. – Vol. 77, № 4. – P. 624–629. DOI: 10.1097/TA.0000000000000406.
129. Comparison of the acute physiology and chronic health evaluation score (APACHE) II with GCS in predicting hospital mortality of neurosurgical intensive care unit patients / A. R. Zali, A. S. Seddighi, A. Seddighi, F. Ashrafi // Global journal of health science. – 2012. – Vol. 4, № 3. – P. 179–184. DOI: 10.5539/gjhs.v4n3p179.
130. Comparison of the Sequential Organ Failure Assessment, Acute Physiology and Chronic Health Evaluation II scoring system, and Trauma and Injury Severity Score method for predicting the outcomes of intensive care unit trauma patients / S. Y. Hwang, J. H. Lee, Y. H. Lee [et al.] // The American journal of emergency medicine. – 2012. – Vol. 30, № 5. – P. 749–753. DOI: 10.1016/j.ajem.2011.05.022.
131. Correlation between arterial blood gas analysis and outcome in patients with severe head trauma / A. Bazzazi, M. A. V. Hasanloei, A. Mahoori [et al.] // Ulusal travma ve acil cerrahi dergisi. – 2014. – Vol. 2, № 4. – P. 236–240. DOI: 10.5505/tjtes.2014.57089.
132. Crash characteristics and pattern of motorcycle related facial bone fractures in a Sub-Urban Nigerian Teaching Hospital / O. S. Obimakinde, O. A. Olajuyin, T. B. Rabi, O. J. Olanrewaju // Nigerian journal of surgery. – 2018. – Vol. 24, № 2. – P. 71–75. DOI: 10.4103/njs.NJS_39_17.

133. Davenport, R. Pathogenesis of acute traumatic coagulopathy / R. Davenport // *Transfusion*. – 2013. – Vol. 53, Suppl. 1. – P. 23S–27S. DOI: 10.1111/trf.12032.
134. del Piñal, F. Severe mutilating injuries to the hand: guidelines for organizing the chaos / F. del Piñal // *Journal of plastic, reconstructive, and aesthetic surgery*. – 2007. – Vol. 60, № 7. – P. 816–827. DOI: 10.1016/j.bjps.2007.02.019.
135. Determination of predictors and risk factors in patients with multiple emergency surgical traumas / S. M. Alamshah, M. Pipelzadeh, S. R. Mousavi [et al.] // *Ulusal travma ve acil cerrahi dergisi*. – 2010. – Vol. 16, № 5. – P. 421–426.
136. Early leukocytosis in trauma patients: what difference does it make? / D. C. Chang, E. E. Cornwell, J. Phillips [et al.] // *Current surgery*. – 2003. – Vol. 60, № 6. – P. 632–635. DOI: 10.1016/j.cursur.2003.07.011.
137. Effects of accidental hypothermia on posttraumatic complications and outcome in multiple trauma patients / P. Mommsen, H. Andruszkow, C. Frömke [et al.] // *Injury*. – 2013. – Vol. 44, № 1. – P. 86–90. DOI: 10.1016/j.injury.2011.10.013.
138. Evolving beyond the vicious triad: Differential mediation of traumatic coagulopathy by injury, shock, and resuscitation / M. E. Kutcher, B. M. Howard, L. L. Sperry [et al.] // *The journal of trauma and acute care surgery*. – 2015. – Vol. 78, № 3. – P. 516–523. DOI: 10.1097/TA.0000000000000545.
139. Facial bone fractures in Ile-Ife, Nigeria: An update on pattern of presentation and care / F. O. Oginni, T. Oladejo, D. P. Alake [et al.] // *Journal of maxillofacial and oral surgery*. – 2016. – Vol. 15, № 2. – P. 184–190. DOI: 10.1007/s12663-015-0826-x.
140. Factors affecting mortality in patients with thorax trauma / S. Emircan, H. Özgüç, S. Akköse Aydın [et al.] // *Ulusal travma ve acil cerrahi dergisi*. – 2011. – Vol. 17, № 4. – P. 329–333.
141. Gastrointestinal function in intensive care patients: terminology, definitions and management. Recommendations of the ESICM Working Group on Abdominal Problems / A. Reintam Blaser, M. L. Malbrain, J. Starkopf [et al.] // *Intensive care medicine*. – 2012. – Vol. 38, № 3. – P. 384–394. DOI: 10.1007/s00134-011-2459-y.

142. Heart rate: is it truly a vital sign? / K. J. Brasel, C. Guse, L. M. Gentilello, R. Nirula // *The Journal of trauma*. – 2007. – Vol. 62, № 4. – P. 812–817. DOI: 10.1097/TA.0b013e31803245a1.
143. Henzler, D. Factors contributing to fatal outcome of traumatic brain injury: a pilot case control study / D. Henzler, D. J. Cooper, K. Mason // *Critical care and resuscitation*. – 2001. – Vol. 3, № 3. – P. 153–157.
144. Hoffman, H. Effect of hypernatremia on outcomes after severe traumatic brain injury: A nationwide inpatient sample analysis / H. Hoffman, M. S. Jalal, L. S. Chin // *World neurosurgery*. – 2018. – Vol. 118. – P. e880–e886. DOI: 10.1016/j.wneu.2018.07.089.
145. Holstein, J. H. What are predictors of mortality in patients with pelvic fractures? / J. H. Holstein, U. Culemann, T. Pohlemann // *Clinical orthopaedics and related research*. – 2012. – Vol. 470, № 8. – P. 2090–2097. DOI: 10.1007/s11999-012-2276-9.
146. Hypernatremia in head-injured patients: friend or foe? / J. F. Payen, P. Bouzat, G. Francony, C. Ichai // *Annales françaises d'anesthésie et de réanimation*. – 2014. – Vol. 33, № 6. – P. 433–435. DOI: 10.1016/j.annfar.2014.05.006.
147. Hypokalemia following trauma / A. L. Beal, K. E. Scheltema, G. J. Beilman, W. E. Deuser // *Shock*. – 2002. – Vol. 18, № 2. – P. 107–110. DOI: 10.1097/00024382-200208000-00002.
148. ICISS: an international classification of disease-9 based injury severity score / T. Osler, R. Rutledge, J. Deis, E. J. Bedrick // *The Journal of trauma*. – 1996. – Vol. 41, № 3. – P. 380–386. DOI: 10.1097/00005373-199609000-00002.
149. Improved predictions from a severity characterization of trauma (ASCOT) over Trauma and Injury Severity Score (TRISS): results of an independent evaluation / H. R. Champion, W. S. Copes, W. J. Sacco [et al.] // *The Journal of trauma*. – 1996. – Vol. 40, № 1. – P. 42–48. DOI: 10.1097/00005373-199601000-00009.
150. Inappropriate prehospital ventilation in severe traumatic brain injury increases in-hospital mortality / T. M. Dumont, A. J. Visioni, A. I. Rughani [et al.] //

- Journal of neurotrauma. – 2010. – Vol. 27, № 7. – P. 1233–1241. DOI: 10.1089/neu.2009.1216.
151. Incidence, etiology, and patterns of maxillofacial fractures in ain-shams university, cairo, egypt: a 4-year retrospective study / A. Mabrouk, H. Helal, A. R. Mohamed, N. Mahmoud // *Craniofacial trauma and reconstruction*. – 2014. – Vol. 7, № 3. – P. 224–232. DOI: 10.1055/s-0034-1374061.
152. Increased nutritional risk in major trauma: correlation with complications and prolonged length of stay / Z. Köseoğlu, M. Ozdoğan, A. Kuvvetli [et al.] // *Ulusal travma ve acil cerrahi dergisi*. – 2011. – Vol. 17, № 6. – P. 521–524. DOI: 10.5505/tjtes.2011.28582.
153. Injury in the United States: 2007 Chartbook / G. Bergen, L. H. Chen, M. Warner, L. A. Fingerhut ; US Department of Health and Human Services, Centres for Disease Control and Prevention, National Centre for Health Statistics. – Hyattsville, 2008. – 179 p.
154. Is the Trauma Mortality Prediction Model (TMPM-ICD-9) a valid predictor of mortality in pediatric trauma patients? / L. D. Cassidy, A. Cook, A. Ertl [et al.] // *Journal of pediatric surgery*. – 2014. – Vol. 49, № 1. – P. 189–192. DOI: 10.1016/j.jpedsurg.2013.09.055.
155. Lemeshow, S. Modeling the severity of illness of ICU patients. A systems update / S. Lemeshow, J. R. Le Gall // *JAMA*. – 1994. – Vol. 272, № 13. – P. 1049–1055.
156. Leukocyte adhesion molecule expression and T cell naïve/memory status following isoproterenol infusion / P. J. Mills, M. Goebel, J. Rehman [et al.] // *Journal of neuroimmunology*. – 2000. – Vol. 102, № 2. – P. 137–144. DOI: 10.1016/s0165-5728(99)00180-0.
157. Major trauma registry of Navarre (Spain): the accuracy of different survival prediction models / T. Belzunegui, C. Gradín, M. Fortún [et al.] // *The American journal of emergency medicine*. – 2013. – Vol. 31, № 9. – P. 1382–1388. DOI: 10.1016/j.ajem.2013.06.026.

158. Management of midface maxillofacial trauma / M. A. Gentile, A. J. Tellington, W. J. Burke, M. S. Jaskolka // Atlas of the oral and maxillofacial surgery clinics of North America. – 2013. – Vol. 21, № 1. – P. 69–95. DOI: 10.1016/j.cxom.2012.12.010.
159. Midfacial fractures: A retrospective etiological study over a 10-year period in Western Romanian population / P. A. Tent, R. I. Juncar, T. Lung, M. Juncar // Nigerian journal of clinical practice. – 2018. – Vol. 21, № 12. – P. 1570–1575. DOI: 10.4103/njcp.njcp_256_18.
160. Morell, V. Predicting severity of trauma by admission white blood cell count, serum potassium level, and arterial pH / V. Morell, E. Lundgren, A. Gillott // Southern medical journal. – 1993. – Vol. 86, № 6. – P. 658–659. DOI: 10.1097/00007611-199306000-00014.
161. Mortality in patients with bilateral femoral fractures / C. E. Copeland, K. A. Mitchell, R. J. Brumback [et al.] // Journal of orthopaedic trauma. – 1998. – Vol. 12, № 5. – P. 315–319. DOI: 10.1097/00005131-199806000-00003.
162. Motamedi, M. H. K. An assessment of maxillofacial fractures: a 5-year study of 237 patients / M. H. K. Motamedi // Journal of oral and maxillofacial surgery. – 2003. – Vol. 61, № 1. – P. 61–64. DOI: 10.1053/joms.2003.50049.
163. Nassar, A. P. Evaluation of Simplified Acute Physiology Score 3 performance: a systematic review of external validation studies / A. P. Nassar, L. M. Malbouisson, R. Moreno // Critical care. – 2014. – Vol. 18, № 3. – Article R117. DOI: 10.1186/cc13911.
164. Neutrophil to lymphocyte ratio as a novel predictor of outcome in patients with severe traumatic brain injury / W. Chen, J. Yang, B. Li [et al.] // The Journal of head trauma rehabilitation. – 2018. – Vol. 33, № 1. – P. E53–E59. DOI: 10.1097/HTR.0000000000000320.
165. Osler, T. A modification of the injury severity score that both improves accuracy and simplifies scoring / T. Osler, S. P. Baker, W. Long // The Journal of trauma. – 1997. – Vol. 43, № 6. – P. 922–926. DOI: 10.1097/00005373-199712000-00009.

166. Paffrath, T. How to define severely injured patients? – An Injury Severity Score (ISS) based approach alone is not sufficient / T. Paffrath, R. Lefering, S. Flohé // *Injury*. – 2014. – Vol. 45, Suppl. 3. – P. S64–S69. DOI: 10.1016/j.injury.2014.08.020.
167. Park, J. M. Neutrophil-to-lymphocyte ratio in trauma patients / J. M. Park // *The journal of trauma and acute care surgery*. – 2017. – Vol. 82, № 1. – P. 225–226. DOI: 10.1097/TA.0000000000001266.
168. Pattern of midface trauma with associated concomitant injuries in a nigerian referral centre / S. Udeabor, V. I. Akinmoladun, A. Olusanya, A. Obiechina // *Nigerian journal of surgery*. – 2014. – Vol. 20, № 1. – P. 26–29. DOI: 10.4103/1117-6806.127105.
169. Pomeranz, S. Hypokalaemia in severe head trauma / S. Pomeranz, S. Constantini, Z. H. Rappaport // *Acta neurochirurgica*. – 1989. – Vol. 97, № 1-2. – P. 62–66. DOI: 10.1007/BF01577741.
170. Poole, G. V. Causes of mortality in patients with pelvic fractures / G. V. Poole, E. F. Ward // *Orthopedics*. – 1994. – Vol. 17, № 8. – P. 691–696. DOI: 10.3928/0147-7447-19940801-07.
171. Prediction of outcome in intensive care unit trauma patients: a multicentre study of Acute Physiology and Chronic Health Evaluation (APACHE), Trauma and Injury Severity Score (TRISS), and 24-hour intensive care unit (ICU) point system / M. J. Vassar, F. R. Jr. Lewis, J. A. Chambers [et al.] // *The Journal of trauma*. – 1999. – Vol. 47, № 2. – P. 324–329. DOI: 10.1097/00005373-199908000-00017.
172. Predictors of mortality following severe pelvic ring fracture: results of a population-based study / B. J. Gabbe, R. de Steiger, M. Esser [et al.] // *Injury*. – 2011. – Vol. 42, № 10. – P. 985–991. DOI: 10.1016/j.injury.2011.06.003.
173. Predictors of poor outcomes after significant chest trauma in multiply injured patients: a retrospective analysis from the German Trauma Registry (Trauma Register DGU®) / S. Huber, P. Biberthaler, P. Delhey [et al.] // *Scandinavian*

- journal of trauma, resuscitation and emergency medicine. – 2014. – Vol. 22, Article 52. DOI: 10.1186/s13049-014-0052-4.
174. Prehospital serum lactate as a predictor of outcomes in trauma patients: a retrospective observational study / F. Guyette, B. Suffoletto, J. L. Castillo [et al.] // *The Journal of trauma*. – 2011. – Vol. 70, № 4. – P. 782–786. DOI: 10.1097/TA.0b013e318210f5c9.
175. Profile, outcome and predictors of mortality of abdomino–pelvic trauma patients in a tertiary intensive care unit in Saudi Arabia / S. H. Haddad, Z. M. Yousef, S. S. Al-Azzam [et al.] // *Injury*. – 2015. – Vol. 46, № 1. – P. 94–99. DOI: 10.1016/j.injury.2014.07.025.
176. Prognostic value of Sequential Organ Failure Assessment and Simplified Acute Physiology II Score compared with trauma scores in the outcome of multiple-trauma patients / P. Fueglistaler, F. Amsler, M. Schüepp [et al.] // *American journal of surgery*. – 2010. – Vol. 200, № 2. – P. 204–214. DOI: 10.1016/j.amjsurg.2009.08.035.
177. Prognostic value of the lethal triad among patients with multiple trauma / M. González Balverde, E. J. Ramírez Lizardo, E. G. Cardona Muñoz [et al.] // *Revista médica de Chile*. – 2013. – Vol. 141, № 11. – P. 1420–1426. DOI: 10.4067/S0034-98872013001100008.
178. Progress in characterizing anatomic injury / W. S. Copes, H. R. Champion, W. J. Sacco [et al.] // *The Journal of trauma*. – 1990. – Vol. 30, № 10. – P. 1200–1207. DOI: 10.1097/00005373-199010000-00003.
179. Propranolol affects stress-induced leukocytosis and cellular adhesion molecule expression / E. C. Kühlwein, M. R. Irwin, M. G. Ziegler [et al.] // *European journal of applied physiology*. – 2001. – Vol. 86, № 2. – P. 135–141. DOI: 10.1007/s00421-001-0526-8.
180. Rating the severity of tissue damage. I. The abbreviated scale / Committee on medical aspects of automotive safety // *JAMA*. – 1971. – Vol. 215, № 2. – P. 277–280. DOI: 10.1001/jama.1971.03180150059012.

181. Red cell distribution width is predictive of mortality in trauma patients / S. Majercik, J. Fox, S. Knight, B. D. Horne // *The journal of trauma and acute care surgery*. – 2013. – Vol. 74, № 4. – P. 1021–1026. DOI: 10.1097/TA.0b013e3182826f02.
182. Risk factors of mortality in severe chest trauma patients / Y. Liu, D. Y. Du, X. Hu [et al.] // *Zhongguo yi xue ke xue yuan xue bao*. – 2013. – Vol. 35, № 1. – P. 74–79. DOI: 10.3881/j.issn.1000-503X.2013.01.014.
183. Rodríguez, M. Predicting mortality from head injury: experience of Sancti Spiritus Province, Cuba / M. Rodríguez // *MEDICC review*. – 2013. – Vol. 15, № 3. – P. 30–33. DOI: 10.37757/MR2013V15.N3.7.
184. Role of oxidative stress in the clinical outcome of patients with multiple blunt trauma / M. T. Gokdemir, O. Sogut, H. Kaya [et al.] // *The Journal of international medical research*. – 2012. – Vol. 40, № 1. – P. 167–173. DOI: 10.1177/147323001204000117.
185. Salentijn, E. G. A ten-year analysis of midfacial fractures / E. G. Salentijn, B. van den Bergh, T. Forouzanfar // *Journal of cranio-maxillofacial surgery*. – 2013. – Vol. 41, № 7. – P. 630–636. DOI: 10.1016/j.jcms.2012.11.043.
186. Santana-Cabrera, L. Differences in the prognosis among severe trauma and medical patients requiring mechanical ventilation / L. Santana-Cabrera, M. Sánchez-Palacios, A. U. Rodríguez // *International journal of burns and trauma*. – 2013. – Vol. 3, № 4. – P. 220–224.
187. Santucci, C. A. Leukocytosis as a predictor of severe injury in blunt trauma / C. A. Santucci, T. B. Purcell, C. Mejia // *The western journal of emergency medicine*. – 2008. – Vol. 9, № 2. – P. 81–85.
188. Serum bicarbonate may replace the arterial base deficit in the trauma intensive care unit / E. FitzSullivan, A. Salim, D. Demetriades [et al.] // *American journal of surgery*. – 2005. – Vol. 190, № 6. – P. 941–946. DOI: 10.1016/j.amjsurg.2005.08.024.
189. Should the IDC-9 Trauma Mortality Prediction Model become the new paradigm for benchmarking trauma outcomes? / A. H. Haider, C. V. Villegas, T. Saleem

- [et al.] // The journal of trauma and acute care surgery. – 2012. – Vol. 72, № 6. – P. 1695–1701. DOI: 10.1097/TA.0b013e318256a010.
190. Søreide, K. Clinical and translational aspects of hypothermia in major trauma patients: from pathophysiology to prevention, prognosis and potential preservation / K. Søreide // Injury. – 2014. – Vol. 45, № 4. – P. 647–654. DOI: 10.1016/j.injury.2012.12.027.
191. Stephenson, S. C. Comparing measures of injury severity for use with large databases / S. C. Stephenson, J. D. Langley, I. D. Civil // The Journal of trauma. – 2002. – Vol. 53, № 2. – P. 326–332. DOI: 10.1097/00005373-200208000-00023.
192. Systematic review of predictive performance of injury severity scoring tools / H. Tohira, I. Jacobs, D. Mountain [et al.] // Scandinavian journal of trauma, resuscitation and emergency medicine. – 2012. – Vol. 20, Article 63. DOI: 10.1186/1757-7241-20-63.
193. Teasdale, G. Assessment of coma and impaired consciousness. A practical scale / G. Teasdale, B. Jennett // Lancet. – 1974. – Vol. 2, № 7872. – P. 81–84. DOI: 10.1016/s0140-6736(74)91639-0.
194. Temiz, A. Admission neutrophil-to-lymphocyte ratio and postoperative mortality in elderly patients with hip fracture / A. Temiz, S. Ersözlü // Ulusal travma ve acil cerrahi dergisi. – 2019. – Vol. 25, № 1. – P. 71–74. DOI: 10.5505/tjtes.2018.94572.
195. Thanapaisal, C. A comparison of the Acute Physiology and Chronic Health Evaluation (APACHE) II score and the Trauma-Injury Severity Score (TRISS) for outcome assessment in Srinagarind Intensive Care Unit trauma patients / C. Thanapaisal, P. Saksaen // Journal of the Medical Association of Thailand. – 2012. – Vol. 95, Suppl. 11. – P. S25–S33.
196. The definition of polytrauma revisited: An international consensus process and proposal of the new 'Berlin definition' / H. C. Pape, R. Lefering, N. Butcher [et al.] // The journal of trauma and acute care surgery. – 2014. – Vol. 77, № 5. – P. 780–786. DOI: 10.1097/TA.0000000000000453.

197. The effect of gender on patients with moderate to severe head injuries / C. Berry, E. J. Ley, A. Tillou [et al.] // *The Journal of trauma*. – 2009. – Vol. 67, № 5. – P. 950–953. DOI: 10.1097/TA.0b013e3181ba3354.
198. The effective risk factors on mortality in patients undergoing damage control surgery / M. Kapan, A. Onder, A. Oguz [et al.] // *European review for medical and pharmacological sciences*. – 2013. – Vol. 17, № 12. – P. 1681–1687.
199. The end of the Injury Severity Score (ISS) and the Trauma and Injury Severity Score (TRISS): ICISS, an International Classification of Diseases, ninth revision-based prediction tool, outperforms both ISS and TRISS as predictors of trauma patient survival, hospital charges, and hospital length of stay / R. Rutledge, T. Osler, S. Emery, S. Kromhout-Schiro // *The Journal of trauma*. – 1998. – Vol. 44, № 1. – P. 41–49. DOI: 10.1097/00005373-199801000-00003.
200. The impact of arterial hypertension on polytrauma and traumatic brain injury / T. Sellmann, D. Miersch, P. Kienbaum [et al.] // *Deutsches Ärzteblatt international*. – 2012. – Vol. 109, № 49. – P. 849–856. DOI: 10.3238/arztebl.2012.0849.
201. The incidence and significance of accidental hypothermia in major trauma – a prospective observational study / S. Ireland, R. Endacott, P. Cameron [et al.] // *Resuscitation*. – 2011. – Vol. 82, № 3. – P. 300–306. DOI: 10.1016/j.resuscitation.2010.10.016.
202. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care / S. P. Baker, B. O’Neill, W. Jr. Haddon, W. B. Long // *The Journal of trauma*. – 1974. – Vol. 14, № 3. – P. 187–196.
203. The neutrophil to lymphocyte ratio on admission and short-term outcomes in orthogeriatric patients / A. Fisher, W. Srikusalanukul, L. Fisher, P. Smith // *International journal of medical sciences*. – 2016. – Vol. 13, № 8. – P. 588–602. DOI: 10.7150/ijms.15445.
204. The predictive value of initial serum lactate in trauma patients / A. Parsikia, K. Bones, M. Kaplan [et al.] // *Shock*. – 2014. – Vol. 42, № 3. – P. 199–204. DOI: 10.1097/SHK.0000000000000208.

205. The relation between the incidence of hypernatremia and mortality in patients with severe traumatic brain injury / U. Maggiore, E. Picetti, E. Antonucci [et al.] // *Critical care*. – 2009. – Vol. 13, № 4. – Article R110. DOI: 10.1186/cc7953.
206. The risk of early mortality of polytrauma patients associated to ISS, NISS, APACHE II values and prothrombin time / L. Mica, K. Rufibach, M. Keel, O. Trentz // *Journal of trauma management and outcomes*. – 2013. – Vol. 7, Article 6. DOI: 10.1186/1752-2897-7-6.
207. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine / J. L. Vincent, R. Moreno, J. Takala [et al.] // *Intensive care medicine*. – 1996. – Vol. 22, № 7. – P. 707–710. DOI: 10.1007/BF01709751.
208. The utility of the predictive scores in polytrauma with abdomino-pelvic injuries: a series of 38 patients / Á. Török, Ş. Bancu, R. Neagoe [et al.] // *Chirurgia*. – 2014. – Vol. 109, № 1. – P. 44–47.
209. Tile, M. Pelvic fractures: operative versus nonoperative treatment / M. Tile // *The Orthopedic clinics of North America*. – 1980. – Vol. 11, № 3. – P. 423–464.
210. Trauma patients with the ‘triad of death’ / B. Mitra, F. Tullio, P. A. Cameron, M. Fitzgerald // *Emergency medicine journal*. – 2012. – Vol. 29, № 8. – P. 622–625. DOI: 10.1136/emj.2011.113167.
211. Trauma score / H. R. Champion, W. J. Sacco, A. J. Carnazzo [et al.] // *Critical care medicine*. – 1981. – Vol. 9, № 9. – P. 672–676. DOI: 10.1097/00003246-198109000-00015.
212. Trauma scores in the management of politrauma patients: which one and what for? / A. N. Sousa, J. A. Paiva, S. A. Fonseca [et al.] // *Acta médica portuguesa*. – 2011. – Vol. 24, № 6. – P. 943–950.
213. Treatment of the severely injured upper extremity / A. Gupta, R. A. Shatford, T. W. Wolff [et al.] // *Instructional course lectures*. – 2000. – Vol. 49. – P. 377–396.

214. Update of the trauma risk adjustment model of the TraumaRegister DGU™: the Revised Injury Severity Classification, version II / R. Lefering, S. Huber-Wagner, U. Nienaber [et al.] // *Critical care*. – 2014. – Vol. 18, № 5. – Article 476. DOI: 10.1186/s13054-014-0476-2.
215. Use of the neutrophil-to-lymphocyte ratio as a component of a score to predict postoperative mortality after surgery for hip fracture in elderly subjects / P. Forget, P. Dillien, H. Engel [et al.] // *BMC research notes*. – 2016. – Vol. 9, Article 284. DOI: 10.1186/s13104-016-2089-0.
216. Validation of the revised injury severity classification score in patients with moderate-to-severe traumatic brain injury / R. Raj, T. Brinck, M. B. Skrifvars [et al.] // *Injury*. – 2015. – Vol. 46, № 1. – P. 86–93. DOI: 10.1016/j.injury.2014.08.026.
217. Vedantam, A. Morbidity and mortality associated with hypernatremia in patients with severe traumatic brain injury / A. Vedantam, C. S. Robertson, S. P. Gopinath // *Neurosurgical focus*. – 2017. – Vol. 43, № 5. – Article E2. DOI: 10.3171/2017.7.FOCUS17418.
218. Victorino, G. P. Does tachycardia correlate with hypotension after trauma? / G. P. Victorino, F. D. Battistella, D. H. Wisner // *Journal of the American College of Surgeons*. – 2003. – Vol. 196, № 5. – P. 679–684. DOI: 10.1016/S1072-7515(03)00128-5.
219. Vrinceanu, D. Principles of surgical treatment in the midface trauma – theory and practice / D. Vrinceanu, B. Banica // *Maedica*. – 2014. – Vol. 9, № 4. – P. 361–366.
220. What is the normal value of the neutrophil-to-lymphocyte ratio? / P. Forget, C. Khalifa, J. P. Defour [et al.] // *BMC research notes*. – 2017. – Vol. 10, № 1. – Article 12. DOI: 10.1186/s13104-016-2335-5.
221. Zahorec, R. Ratio of neutrophil to lymphocyte counts—rapid and simple parameter of systemic inflammation and stress in critically ill / R. Zahorec // *Bratislavské lekárske listy*. – 2001. – Vol. 102, № 1. – P. 5–14.

222. Zandi, M. The relationship between head injury and facial trauma: a case-control study / M. Zandi, S. R. Seyed Hoseini // Oral and maxillofacial surgery. – 2013. – Vol. 17, № 3. – P. 201–207. DOI: 10.1007/s10006-012-0368-z.